## Biological and fisheries data for managing the deep-sea crabs *Hypothalassia acerba* and *Chaceon bicolor* in Western Australia

Smith, K.D., Potter, I.C. and Hall, N.G.





**Australian Government** 

Fisheries Research and Development Corporation

# Projects No. 1999/154 and 2001/055

Fisheries Research and Development Corporation Report FRDC projects 1999/154 and 2001/055 Fisheries Research and Development Corporation Report FRDC projects 1999/154 and 2001/055

## FINAL REPORT

## Biological and fisheries data for managing the deep-sea crabs *Hypothalassia acerba* and *Chaceon bicolor* in Western Australia

Smith, K. D. Potter, I. C. Hall, N. G.

September 2004

ISBN: 0-86905-883-5

Centre for Fish and Fisheries Research Murdoch University Murdoch, Western Australia 6150

This work is copyright. Except as permitted under the Copyright Act 1968 (Cth), no part of this publication may be reproduced by any process, electronic or otherwise, without the specific written permission of the copyright owners. Neither may information be stored electronically in any form whatsoever without such permission.

### **Table of Contents**

NON-TECHNICAL SUMMARY:			
AC	KNOWL	EDGEMENTS	. 10
1.0	GEN	NERAL INTRODUCTION	.11
1.1	THE C	CHAMPAGNE CRAB HYPOTHALASSIA ACERBA	. 11
1.2	THE C	CRYSTAL CRAB CHACEON BICOLOR	. 11
1.3	PREV	IOUS BIOLOGICAL STUDIES ON DEEP-SEA CRABS	. 13
1.4	NEED	)	. 14
2.0	GEN	NERAL MATERIALS AND METHODS	. 16
2.1	SAMP	LING OF HYPOTHALASSIA ACERBA	. 16
	2.1.1	Depth-stratified sampling regime	. 16
	2.1.2	Onboard sampling of commercial catches of Hypothalassia acerba	. 22
	2.1.3	Structure of traps used in the commercial fishery to catch Hypothalassia acer	ba.
	214	Tagging of Hypothalassia acerba	. 22
	2.1.5	Sampling of Chaceon bicolor	.23
	2.1.6	Structure of traps used in the commercial fishery to catch Chaceon bicolor	. 24
2.2	MEAS	SUREMENTS	. 25
	2.2.1	Water temperatures	. 25
	2.2.2	Data recorded for crabs	. 25
3.0	THE	E FISHERIES FOR HYPOTHALASSIA ACERBA AND CHACEON	
BIC	COLOR		. 29
3.1	INTRO	DDUCTION	. 29
3.2	MATE	ERIALS AND METHODS	. 30
	3.2.1	Data on commercial catches	. 30
	3.2.2	Onboard collection of data	. 30
	Chaceon	bicolor	.31
33	J.Z.J RESU	Dala Analysis	. 32
5.5	331	Commercial catch and effort data for Hypothalassia acerba	. 33
	3.3.2	Commercial catch and effort data for Chaceon bicolor	. 41
	3.3.3	Analysis of catches and sizes of crabs derived from onboard sampling	. 41
	3.3.4	Carapace Lengths	. 49
	3.3.5	Carapace lengths of Chaceon bicolor in non-meshed and meshed traps	. 53
	3.3.6	Results from release of tagged Hypothalassia acerba	. 54
3.4	DISCU	JSSION	. 54
	3.4.1	Annual trends in catch statistics	. 54
	3.4.2	Monthly trends in catch statistics	.57
	3.4.3	Calcres and carapace lengths of Hypothalassia acerba and Chaceon bicolor	. 38
4.0	REL	ATIVE ABUNDANCES AND SIZE COMPOSITIONS OF	
	POTHALA DTUG AN	ASSIA ACERBA ON TWO COASTS AND IN DIFFERENT WATER	50
DEI			. 37
4.1	INTRO	ODUCTION	. 59

4.2	MAT	ERIALS AND METHODS	61
4.3	RESU	ILTS	64
	4.3.1	Water temperature	64
	4.3.2	Relative abundance of crabs in different water depths and temperature	64
	4.3.3	Relationship between carapace length and water depth, coast and season	69
	4.3.4	Sex ratios	74
	4.3.5	Results of enclosing traps with mesh	75
4.4	DISC	USSION	75
	4.4.1	Relationship between catch rates and water depth	75
	4.4.2	Relationship between catch rates and water temperature	79
	4.4.3	Trap selectivity	80
5.0	CO	MPARISONS BETWEEN THE REPRODUCTIVE BIOLOGY OF	
FEI	MALES (	OF TWO SPECIES OF DEEP SEA CRABS WHICH LIVE IN DIFFERI	ENT
WA	TER DE	PTHS	81
51	INTR	ODUCTION	81
5.1	MAT	FRIALS AND METHODS	01 83
5.2	521	Collection of samples	05 83
	5 2 2	Laboratory processing	05
53	D.2.2 RESI	IT TS	05 85
5.5	5 3 1	Ovarian maturation stages	05 85
	532	Monthly trands in reproductive indices	80 80
5 /	DISC	USSION	09 08
J. <del>T</del>	5 1 1	Comparisons between Hypothalassia acerba on the west and south coasts	00 08
	5.4.2	Comparisons between Hypothalassia acerba and Chaceon bicolor	101
	5.4.3	Investment in gonadal development by Hypothalassia acerba	102
<i>c</i> 0	5.1.5 DOI		. 102
6.0	PO'	TENTIAL BIAS IN ESTIMATES OF THE SIZE OF MATURITY OF	104
CK	ABS DEI	RIVED FROM TRAP SAMPLES	104
6.1	INTR	ODUCTION	104
6.2	MAT	ERIAL AND METHODS	106
	6.2.1	Sampling and measurement of crabs	106
	6.2.2	Data analysis	. 108
6.3	RESU	/LTS	108
	6.3.1	Size composition	108
	6.3.2	Size at maturity	110
6.4	DISC	USSION	114
7.0	DO	THE CHELAE OF THE MALES OF THREE CRAB SPECIES UNDER	RGO
AL	LOMETI	RIC CHANGES THAT CAN BE USED TO DETERMINE	
MC	RPHOM	ETRIC MATURITY?	117
71	INTR	ODUCTION	117
7.1	MAT	FRIAL AND METHODS	120
1.2	721	Collection of crabs	120 120
	7.2.1	Measurements	120
	722	Relationship between propodus length and body size	120
	72.5	Probability of selecting an incorrect model	121
	725	Physiological maturity	124
	726	Size at physiological maturity	125
73	RESE	II TS	126

	7.3.1	Relationship between propodus length and body size	
	7.3.2	The probability of selecting an incorrect model form	
	7.3.3	Size at physiological maturity	
7.4	DISC	USSION	
	7.4.1	Morphometric and physiological maturity of Portunus pelagicus	
	7.4.2	Morphometric and physiological maturity of Hypothalassia acerba a	and Chaceon
	bicolor		
8.0	REI	FERENCES	135
9.0	BEN	NEFITS	144
10.0	FUI	RTHER DEVELOPMENT	144
11.0	PLA	NNED OUTCOMES	145
12.0	GEI	NERAL CONCLUSIONS	147
13.0	API	PENDICES	151

2001/055	Biological and fisheries data for managing deep sea crabs in Western
	Australia

PRINCIPAL INVESTIGATOR:	Professor I. C. Potter
ADDRESS:	Centre for Fish and Fisheries Research Division of Science and Engineering Murdoch University South Street, Murdoch WA 6150 Ph: 08 9360 2524 Fax: 08 9360 6303

#### **OBJECTIVES:**

#### The champagne crab Hypothalassia acerba (FRDC 1999/154)

- 1. Determine the composition by sex, size (carapace length (CL), to the nearest 1 mm), and weight (to the nearest 1 g) of champagne crabs in commercial crab pots from different depths in the two regions in which the champagne crab is fished , *i.e.* Geraldton to Fremantle and Augusta to Albany.
- 2. Use traps with a small mesh size to determine the full range in size (CL) and weight of crabs in different depths in the two sampling regions.
- 3. Produce preliminary estimates of growth of female and male crabs, based on growth increments and inter-moult periods.
- 4. Obtain indications of the movements of crabs and their relationships to body size and reproductive status.
- 5. Determine the carapace lengths at which 5 and 50% of female and male crabs reach maturity.
- 6. Determine the duration and location (area and water depth) of spawning and the fecundity and its relationship to body size.
- 7. Determine the diets of crabs and how they vary with body size and shell state. N.B. This study, which will also provide information on the type of habitat occupied by champagne crabs, will be carried out by a student at no cost to the project.
- 8. Establish the distribution patterns, size composition and prevalence of ovigerous females, and analyse data for other biological variables that were observed by fishers and recorded in log books.
- 9. The yield and egg (or mature biomass) per recruit relationships.
- 10. Assess the management implications of the biological data, and provide the results to managers in a form that can be used to help manage and conserve the fishery for the champagne crab.

#### The crystal crab Chaceon bicolor (FRDC 2001/055)

- 1. Implement a voluntary log book system in which fishers will record the number, size and location of capture of crabs and establish the data bases that are necessary to record and maintain logbook, size composition and tagging data.
- 2. Collect and process biological data on the size, weight, moult stage, shell state, sex ratio, gonadal weight and gonadal stage of crystal crabs caught by commercial fishers at regular intervals throughout the year.

- 3. Determine the relative abundance of crystal crabs in the various depths and regions in which this species is fished.
- 4. Determine the pattern of movement, growth and frequency of spawning of crystal crabs.
- 5. Estimate the carapace lengths at which 5 and 50% of female and male crabs reach maturity.
- 6. Determine the duration and location (area and water depth) of spawning and the fecundity and its relationship to body size.
- 7. Determine the size of the gaps that are required in the traps to allow undersize crabs to escape.
- 8. Determine the distribution patterns, size composition and the prevalence of ovigerous females, and other biological variables, observed by fishers and recorded in log books.
- 9. Undertake a depletion study to provide direct estimates of catchability and densities in different area.
- 10. Provide an initial assessment of the status of the crystal crab stock on the west coast of Western Australia.

Note that 1,4,7,8,9 and 10 will be undertaken by the Department of Fisheries Western Australia as its contribution to FRDC Project No. 2001/055 and the results of those components will be provided in a separate final report produced by the Department when the project finishes on 30 June 2005.

#### **NON-TECHNICAL SUMMARY:**

#### OUTCOMES ACHIEVED TO DATE

Biological data for the champagne crab *Hypothalassia acerba* and crystal crab *Chaceon bicolor* have been collected to enable managers to develop appropriate plans for conserving the stocks of these two commercial species of deep sea crabs. Emphasis was placed on outlining the catch statistics for the early years of the fisheries and on determining the depth distributions, size compositions, sizes at maturity, spawning periods, sex ratios and fecundity of these species. The use of samples of female champagne and crystal crabs collected by traps (their only method of capture) almost certainly leads to the size at gonadal maturity of those crabs to be underestimated. This needs to be taken into account if size at maturity is to be used for setting minimum legal lengths. Since the patterns of growth of the claws of male champagne and crystal crabs do not change at the pubertal moult, they could not be used in the traditional manner to determine maturity. The implications of these and our other results have been discussed with Dr Roy Melville Smith (WA Fisheries), who is continuing research into the biology of the crystal crab.

During recent years, commercial trap fisheries have developed for the champagne and crystal crabs in deep waters off the lower west and south coasts of Western Australia. Thus, data on crucial aspects of the biology of these species were required to enable plans to be developed for conserving their stocks. No such biological data were previously available.

Commercial catches of champagne crabs in south-western Australia rose from negligible levels in 1990 to peak at ~73 000 kg in 1998 and then declined progressively to ~17 300 kg in 2003, due mainly to fishers on the west coast shifting their effort towards crystal crabs. The catches of champagne crabs underwent a progressive seasonal change, particularly on the south coast where they reflected a decline in fishing effort during the inclement late autumn to middle spring months. Commercial catches of crystal crabs in south-western Australia, which came almost entirely from the lower coast, rose from very low levels in 1997 to reach ~222 000 kg in 2001 and then remained close to this level in 2002 and 2003. Catches and catch per unit effort of crystal crabs remained similar throughout the year. For the fisheries for both champagne and crystal crabs, the mean carapace lengths and catch rates were greater for males than females. Enclosure of traps in fine mesh did not lead to the capture of a relatively greater number of small crabs.

On the west and south coasts, the abundance of champagne crabs peaked at depths of 200 and 145 m, respectively, but at similar temperatures of 16.1-17.1°C. On the former coast, the crystal crab lives in depths of 450 to 1 220 m, where temperatures are only 4 to 6.5°C. The percentage of males in catches of champagne crabs was less on the west (69%) than south coast (84%). The carapace length of champagne crabs declined significantly by 4 mm with each 100 m increase in depth. As with commercial catches, the maximum and mean carapace lengths for males were greater than for females on both the west and south coasts. From the above, it follows that, in the case of champagne crabs, (1) distribution is related to water depth and temperature, (2) body size is inversely related to water depth and (3) males grow larger and are caught in greater numbers than females. There was also evidence that the distribution of champagne crabs changed slightly with season and that champagne crabs and other large deep water invertebrate predators tended to live in different water depths and thus avoid major competition for resources.

The trends exhibited by reproductive variables demonstrate that champagne crabs reproduce seasonally on the lower west coast, with females maturing progressively between July and December and becoming berried (egg bearing) between January and March. The reproductive characteristics of champagne crabs on the south coast differed markedly, in the following ways, from those on the lower west coast. (i) No berried females and only two females with egg remnants were caught. (ii) The development of large mature eggs was delayed until females reached a larger size. (iii) Investment of food resources in gonadal development was less. These results strongly indicate that (i) conditions on the south coast are not as conducive for ovarian development and reproduction and (ii) females migrate from the south to lower west coast for spawning. In contrast to champagne crabs, crystal crabs reproduce throughout all or most of the year on the lower west coast, presumably reflecting their occupancy of far deeper waters where environmental conditions vary less during the year. Although the mean weights of berried females of champagne and crystal crabs were not significantly different (P > 0.05), the mean fecundity of the former species (356 210) was significantly greater (P < 0.001) than that of the latter species (192 070). The high fecundity of the champagne crab may reflect adaptations by this seasonal spawner to optimise egg production during its relatively short breeding season.

The size at onset of sexual maturity (SOM) of female crustaceans is typically estimated by analysing the proportions of mature females in sequential size classes. The validity of this approach depends on the composition of the samples reflecting accurately that present in the environment. However, catches obtained by traps, a passive fishing method, typically contain a disproportionately greater numbers of large crabs, whereas those obtained using active fishing methods, such as seine netting and otter trawling, represent far better the size composition of the population. From data collected during a previous study, the samples of female blue swimmer crabs, *Portunus pelagicus*, caught by trapping were predominantly mature, whereas those collected by seining and trawling contained numerous immature as well as mature females. Thus, the samples of females collected by trap are biased towards mature crabs, which results in an underestimate of the SOM. From the results for the blue swimmer crab, the mature females of the champagne and crystal crab would also be likely to particularly susceptible to capture by traps, their only current method of capture. The respective sizes at which, on average, the females of these two species reach maturity, based on trap samples, namely carapace lengths of 69.7 and 90.5 mm, are thus likely to be lower than would be the case with truly representative samples.

The fact that the claws of male crabs are frequently assumed to undergo a change in relative size at the pubertal moult has been used as the basis for determining the size of male crabs at maturity. Initial plots of claw length versus carapace width (CW) or length (CL) of the males of blue swimmer, champagne and crystal crabs revealed no conspicuous change in the relative sizes of the claws. However, detailed statistical analysis revealed that the claws of blue swimmer crabs do undergo a change in the pattern of growth at 80 mm CW, namely at 8 mm less than their width at gonadal maturity. In contrast, no such evidence was found that the claws of either champagne or crystal crabs undergo similar changes. Maturity could not therefore be determined for these two species from the relative lengths of their claws. The lengths at gonadal maturity of trap-caught males of champagne and crystal crabs were 68.1 and 94.3 mm CL, respectively.

It follows that the establishment of minimum legal sizes for the males and females of both champagne and crystal crabs has to be based on the lengths at which gonadal maturity is attained, but recognising that, in the case of females, these values are likely to underestimate slightly the true average size at maturity. Size-frequency data indicate that, by the time champagne and crystal crabs are caught in traps, they are approaching their maximum sizes. Thus, if fishing effort increases, yield per recruit will increase and spawning biomass per recruit will decrease. Consequently, exploitation should be controlled to ensure that, in the future, catch per unit of effort is maintained above a specified minimum acceptable level.

**KEYWORDS:** crabs, deep sea, distribution, depth, temperature, size, reproduction, maturity

#### ACKNOWLEDGEMENTS

We are very grateful to the deep-sea crab fishers Tim Goodall, David and Lance Hand, Cliff Neave, Graham Pateman and Gavin Wilson for their invaluable assistance with sampling crabs and to Kevin Eiden, Brenden Muguire and Ray Prior for the use of their vessels. We also welcomed the constructive advice of Andrew Levings at the initial stages of the project and the invaluable discussions held with Roy Melville Smith during this study. Sincere appreciation is extended to Simon de Lestang, Duncan Forbes, Simon Hambleton, Alex Hesp, Rick Hoddell, Matt Hourston, Stephan Linke, Chris Marsh, Matt Pember, Norman Ramsay, Erug Sezmis, Justin Smith, Michael Travers, Ayesha Whitehead and Brenden Worth who helped with the field work which was often conducted in inhospitable conditions. Finally, special thanks are extended to Mark Cliff for his patience and efforts in extracting catch statistics from files at the Western Australian Department of Fisheries.

#### **1.0 GENERAL INTRODUCTION**

#### 1.1 THE CHAMPAGNE CRAB HYPOTHALASSIA ACERBA

*Hypothalassia acerba* Koh and Ng (**Figure 1.1a**), which used to be referred to as the spiny crab but is now termed the champagne crab, belongs to the Family Eriphiidae. This species lives in deep water off the south-western Australian coast and has recently been recognised as distinct from the only other member and type species of the genus *Hypothalassia*, namely *Hypothalassia armata*, which is restricted to the Pacific (Koh and Ng, 2000). Commercial catches indicate that, in Western Australia, the distribution of *H. acerba* extends from at least as far north as Kalbarri at ~114°E 27°S on the west coast, southwards and eastwards to at least Eucla at ~129° E, 32° S on the south coast.

Until 1989, the champagne crab was only exploited on the west coast of Australia by western rock lobster fishers working in deep water during December and January. However, since 1989, a small number of fishers have focused on catching *H. acerba* on this coast under a deep-sea crab endorsement license. On the south coast, champagne crabs have formed, since 1990, a component of a multi-species fishery, which also includes the king crab *Pseudocarcinus gigas* and the southern rock lobster *Jasus edwardsii*. A live export fishery for *H. acerba* commenced in 1989 and the Western Australian Fisheries Department Catch and Effort Statistics (CAES) show that, during the next decade, catches (total live weight) rose progressively to reach a peak of over 75 000 kg in 1998.

#### 1.2 THE CRYSTAL CRAB CHACEON BICOLOR

*Chaceon bicolor* Manning and Holthuis (**Figure 1.1a**), which used to be referred to as the snow crab but is now termed the crystal crab, belongs to the family Geryonidae. This species is found in deep waters off large regions of the Australian and New Zealand coastlines (Jones



<image>

**Figure 1.1** (a) Male *Hypothalassia acerba* (above) and male *Chaceon bicolor* (below) and (b) the fishing vessel *Barbarossa* which was used on the lower west coast for the depth-stratified sampling regime (see Chapter 2).

and Morgan 1996). A live export fishery for *C. bicolor* from Western Australia commenced in 2000 and catches increased to 143 000 kg during 2000 (Western Australian Department of Fisheries CAES).

A comprehensive account of the fisheries for both *H. acerba* and *C. bicolor* in Western Australia is provided in Chapter 3.

#### 1.3 PREVIOUS BIOLOGICAL STUDIES ON DEEP-SEA CRABS

Prior to the present study, no biological data of the type required for fisheries management had been collected for either *H. acerba* or *C. bicolor*. However, the biology of *Pseudocarcinus gigas*, which is the most valuable of Australia's deep-water crabs and belongs to the same family (Eriphiidae) as *H. acerba*, have been studied (Gardner, 1996; Levings *et al.*, 1996; Gardner, 1997; Gardner and Northam, 1997; Gardner, 1998a,b; Levings *et al.*, 2001; Gardner and Williams, 2002; McGarvey *et al.*, 2002). Thus, there are now data on the following aspects of *P. gigas*, namely geographical distribution, growth, reproductive biology, egg production and size at sexual maturity and on fisheries aspects, such as catches and catch per unit effort, the water depths at which king crabs are caught, and the size structure of the catches. Furthermore, the larvae of *P. gigas* have been reared in the laboratory to the megalopae stage (Gardner, 1996; Gardner and Northam, 1997).

Established deep-sea crab fisheries exist for the geryonid species, *Chaceon maritae*, off south west Africa (Melville Smith, 1988) and *Chaceon quinquedens* off the southern coast of New England (Wigley *et al.*, 1975). Smaller scale exploratory fisheries have also existed over the years for *Chaceon affinis* off Portugal (Pinho *et al.*, 2001), the Canary islands (López Abellán *et al.*, 2002) and in the Gulf of Mexico (Haefner, 1978) and for *C. quinquedens* in the Gulf of Mexico (Haefner, 1978). Several species of deep-sea crab are exploited off Alaska, including the lithodid *Paralithodes camtschaticus* (Zhou and Shirley, 1997), and the majids *Chionoecetes tanneri*, *C. bairdii* and *C. opilio* (Claxton and Govind, 1993). *Chionoecetes* 

*opilio* is also fished off the Atlantic coast of Canada (Sainte Marie *et al.*, 1997). *Chionoecetes opilio*, *C. japonicus* and *C. bairdii* are fished off Japan (Kon, 1996).

#### 1.4 NEED

The successful management of a fishery for any species is dependent on fisheries managers having reliable data on certain key aspects of the biology of that species. The growth rate, fecundity and frequency of spawning are of particular importance as they are essential for calculating yield and egg (or mature biomass) per recruit relationships, which may be used to establish an optimal legal minimum size at first harvesting or for assessment of the status of the stock. The sizes at which 5 and 50% of crabs reach maturity provide a convenient description of the logistic relationship between the proportion of crabs that are mature and body size, when this relationship is used for setting an appropriate minimum legal size at capture. The pattern of movements and the way in which they are related to size and reproductive condition are important for interpreting catch and effort data. No data are available for any of the above aspects of the biology of either the champagne crab *Hypothalassia acerba* or the crystal crab *Chaceon bicolor*.

The champagne crab and crystal crab are assumed to be slow growing and long-lived and therefore vulnerable to overfishing. For that reason, the massive increase that occurred in the catches of these species since they started to be fished commercially provided an urgent need for appropriate management arrangements to be set in place to protect the stocks of these brachyurans. The biological information required for developing management plans for these species include growth rates, size composition, size at first maturity, spawning period, frequency of spawning, fecundity and yield and egg (or mature biomass) per recruit relationships. Furthermore, there was an urgent need to collect data on the catch statistics, size compositions, sex ratios and relative abundances by depths of *H. acerba* and *C. bicolor* in the early stages of the development of the fisheries for these two species, so that any changes in these important variables in the future could be detected.

Thus, there was a pressing need to collect baseline biological and fishery data, as listed under objectives, which could be used to produce a preliminary stock assessment for assisting in the management and conservation of these species in Western Australia. Such a stock assessment was required to meet Environment Australia's Schedule 4 exemption in the short term, and Fisheries WA/Environment Australia's requirements which were to be initiated in December 2002.

#### 2.0 GENERAL MATERIALS AND METHODS

#### 2.1 SAMPLING OF HYPOTHALASSIA ACERBA

The biological data reported for champagne crabs in Chapters 4 to 7 were obtained from three sampling regimes. The first sampling regime was designed specifically to obtain monthly data on the relative abundance, reproductive biology and size composition of *H. acerba* in different water depths off the lower west and south coasts of Western Australia (Chapter 4). It is subsequently referred to as the depth-stratified sampling regime to distinguish it from the second sampling regime which involved obtaining data for *H. acerba* by examining crabs while onboard commercial fishing vessels during their normal fishing activities. The third sampling regime involved collecting or purchasing samples of commercial catches from fish markets.

#### 2.1.1 Depth-stratified sampling regime

For the depth-stratified sampling regime, *Hypothalassia acerba* was sampled off the lower west coast of Western Australia at a latitude of  $\sim 30^{\circ}45$ /S and a longitude of  $\sim 114^{\circ}50$ /E and off its southern coast at a latitude of  $\sim 35^{\circ}15$ /S and a longitude of  $\sim 118^{\circ}30$ /E (**Figure 2.1**). Seven sites (water depths) were sampled on each sampling occasion on both coasts. The sites were selected so that they were located in areas where the mean water depths were 35, 90, 145, 200, 255, 310 and 365 m, respectively, *i.e.* sites were equally spaced at depth intervals of 55 m. They were located at distances from the shore of 21.6, 28.5, 34.1, 40.1, 40.8, 40.9 and 41.1 km off the lower west coast and of 0.5, 28.2, 31.8, 32.1, 32.5, 32.7 and 33.1 km from the shore off the south coast.

The seven sites on the lower west coast and seven sites on the south coast were sampled in nine and four months, respectively, between January 2000 and April 2002 with each coast



**Figure 2.1.** Map showing the region of south-western Australia where *Hypothalassia acerba* and *Chaceon bicolor* were caught. Boxes show location of the depth-related sampling regime on the west and south coasts. Circles designate the location where the samples at the different depths were collected. Each sampling unit contained one rectangular trap at each end and a central circular trap (see bottom left hand corner).

being sampled at least once in each calendar season (**Table 2.1**). Samples were collected from each site (water depth) on each sampling occasion using three lines of traps, with each line of 11 mm propylene rope having a rectangular trap at each end and a circular trap in the middle (see **Figure 2.2a**). The rectangular traps had a steel base and timber frame that was 960 mm long x 800 mm wide x 460 mm high and contained an entrance 270 mm in diameter x 170 mm in depth at the top. The trap bases consisted of parallel steel rods, 5 mm in diameter, situated 80 mm apart. The sides, top and ends of the traps were enclosed with wooden battens approximately 70 mm apart. The round traps were 880 mm in diameter x 450 mm in height and contained an entrance of 280 mm diameter x 200 mm deep at the top. Both of the trap types described above, which are typically employed to catch the western rock lobster *Panulirus cygnus* (George, 1962), are the same as those used by commercial fishers to catch champagne crabs.

In each water depth, the length of rope between the trap and the water surface was always approximately twice the water depth. A distance of 72 m separated the traps in each line (**Figure 2.2b**). Each line was set ~500 m apart. Escape gaps, a legal requirement of these commercial western rock lobster traps, were covered. Traps were baited with fish heads, such as those of the orange roughy *Hoplostethus atlanticus* Collett or the New Zealand hoki *Macryronus novaezelandiae* (Hector) which are commonly used to catch these crabs, and also with cattle hide or foreleg, which is more resistant to lice (isopods and amphipods) than the fish heads and thus last longer. A single inflatable fender and up to 5 cast alloy buoys were used to keep the surface rigs afloat (**Figure 2.3a**). Traps were typically set for five days.

In an attempt to catch greater numbers of small crabs off the lower west coast during May (autumn) and November (spring) 2001, the rectangular traps at each end of an additional equal number of lines were enclosed with welded galvanised aviary mesh with meshes of 12.7  $\times$  25.4 mm (**Figure 2.3b**). The mesh, which was attached to the traps using cable ties, nails



**Figure 2.2.** (a) Round cane stick traps (above) and rectangular jarrah slat traps (below) used for the depth - stratified sampling regime and (b) warning ropes (blue) to which the traps were attached. The traps were separated by 72 m of ground rope (yellow).



**Figure 2.3.** (a) Inflatable buoy, one of which was attached to each line of traps and (b) rectangular jarrah trap enclosed with wire mesh.

and staples, enclosed the whole of the traps, except for their entrance. Since the spaces between the canes used in the central circular trap were small, it was not considered necessary also to enclose these traps in aviary mesh. The additional lines of small mesh traps were used to fish the same water depths, but  $\sim$ 2 km further south, effectively replicating the experiment in those two months. In all of the sampling conducted off the lower west coast after November 2001, the rectangular traps were enclosed with aviary mesh (see **Table 2.1**).

**Table 2.1**Dates and seasons when traps were used to catch *Hypothalassia acerba* off thelower west and south coasts of Western Australia.n.c., not covered with aviary mesh; c,covered with aviary mesh.

Date	Season		
West coast	West coast		
17.02.00	Summer, n.c.		
23.8.00	Winter, n.c.		
3.3.01	Autumn, n.c.		
18.9.00	Spring, n.c.		
12.5.01	Autumn , n.c. & c		
6.11.01	Spring, n.c. & c		
18.12.01	Summer, c		
20.1.02	Summer, c		
25.4.02	Autumn, c		
South coast			
19.3.00	Autumn, n.c.		
17.6.00	Winter, n.c.		
25.11.00	Spring, n.c.		
22.2.02	Summer, n.c.		

#### 2.1.2 Onboard sampling of commercial catches of Hypothalassia acerba

Samples of *Hypothalassia acerba* were obtained onboard from the trap catches of commercial fishers operating off the lower west and south coasts of Western Australia between ~30° 10'S, 114° 30'E (~ west of Jurien) and 32° 30'S, 114° 60'E (west of Mandurah) and between ~34° 40'S, 115° 40'E (south of Augusta) and 34° 45'S, 119° 30'E (south of Bremer Bay), respectively (**Figure 2.1**). Off the lower west and south coasts, the water depths fished by commercial fishers during these monitoring cruises, ranged from 180 to 280 m and 70 to 240 m respectively. These trips were undertaken at regular intervals off the lower west coast between July 1999 and December 2000 and off the south coast between July 1999 and February 2002, respectively.

#### 2.1.3 Structure of traps used in the commercial fishery to catch Hypothalassia acerba

The single commercial fisher, who targeted champagne crabs on the west coast, consistently used rectangular jarrah rock lobster traps of the type employed in the depth-stratified sampling regime described in Section 2.1.1. Note that other fishers, who were targeting western rock lobster on the west coast, were not allowed to retain for sale any incidental catches of champagne crabs.

Since there is no legislation specifying that a particular type of trap should be used by commercial fishers for catching champagne crabs, the trap types used on the south coast varied. Note that, on the south coast as opposed to the west coast, some fishers possessed licenses that enabled them to sell any catches of *H. acerba* that were obtained incidentally when they were fishing for other species such as the southern rock lobster *Jasus edwardsii* and the king crab *Pseudocarcinus gigas*.

The three main types of trap used on the south coast included the circular cane stick traps of the type used in the depth-stratified sampling regime (Section 2.1.1). The first of the

other two trap types was round-sided with flat tops (1400 mm base diameter, 350 mm high, 272 mm neck diameter), while the second was conical and beehive shaped (860 mm base diameter, 407 mm high, 272 mm neck diameter). Both of these latter traps were enclosed with either diamond-shaped stainless steel chicken wire or resilient synthetic netting. All three traps were weighted with ~ 40 kg of steel ballast.

For commercial fishing off both coasts, between two and twelve traps were strung together in lines, with varying distances of rope between the traps. Furthermore, soak time ranged from four days to several weeks, mainly due to variations in weather conditions. Traps were usually baited with fresh or salted heads of fish species such as the orange roughy *Hoplostethus atlanticus* Collett or the New Zealand hoki *Coryphaenoides novaezelandiae* Hector. Cattle hide or foreleg was employed until its use was banned in January 2002.

#### 2.1.4 Tagging of Hypothalassia acerba

Crabs were tagged using Hallprint TBA-1 t-bar tags. The tags were inserted using a Paxar tagging gun on the right hand side medial epimeral suture line of the carapace, between the fourth and fifth coxa, with the t-bar lodged in the branchial cavity (**Figure 2.4a**). After tagging, crabs were returned immediately to the water in order to minimise their displacement due to drift. Crabs with missing appendages were not tagged.

#### 2.1.5 Sampling of Chaceon bicolor

Sampling for *C. bicolor* aboard commercial fishing vessels was undertaken on four occasions during 2000 and, wherever possible, bimonthly between June 2001 and December 2003.



**Figure 2.4.** (a) Female *Hypothalassia acerba* showing the t-bar anchor tag inserted within the epimeral suture line of the carapace and (b) commercial fishing trap used to catch *Chaceon bicolor*. (Photo courtesy of S. Norton).

#### 2.1.6 Structure of traps used in the commercial fishery to catch Chaceon bicolor

*Chaceon bicolor* is fished commercially using lines containing up to 100 plastic recreational western rock lobster traps, which are 675 mm long, 350 mm wide and 475 mm high and possess, at their top, an entrance of 170 mm diameter and 200 mm depth (**Figure 2.4b**). The traps are made of light-weight plastic to prevent them sinking into the soft substrate and are baited with heads of the New Zealand hoki or whole Australian salmon, *Arripis truttaceus* (Cuvier).

#### 2.2 MEASUREMENTS

#### 2.2.1 Water temperatures

On each occasion that the depth-stratified sampling for *H. acerba* was implemented, the bottom water temperature at each site was recorded using StowAway TidbiT HTI-537 digital temperature loggers that were enclosed in protective wire mesh and attached to the inside top of one of the traps. As a logger very occasionally malfunctioned, the temperature in that depth was not recorded on that sampling occasion. Bottom water temperature was recorded using the same devices, which were attached to commercial traps deployed to catch *C. bicolor*. The temperature was recorded to 0.1°C at 5 min intervals and the resultant values downloaded from these loggers to a computer and subsequently averaged for each sampling occasion at each sampling site.

#### 2.2.2 Data recorded for crabs

In the field, the contents of each individual trap were recorded separately and a note made of the location, date and depth (m) at which all crabs were caught. The following characteristics of each crab were recorded. (1) The carapace length (*CL*) of each *H. acerba* and *C. bicolor*, *i.e.* the distance from the midpoint between the bases of the two anterior medial horns and the

posterior margin of the carapace, was measured to the nearest 1 mm. This is the typical way of measuring body size in deep-water crabs (Levings et al., 1996; Gardner, 1997; Goshima and Kanazawa, 2000). It is preferable to using carapace width, *i.e.* the distance between the two lateral spines of the carapace, as those structures are particularly prone to wear. (2) In the case of females, pleopod condition was recorded as either clean with no sign of egg attachment, or as having either eggs (berried) or egg remnants attached. (3) For each female, a record was kept of the relative size and shape of the gonopores and whether mating abrasions were present around their gonopores. Most females of C. bicolor with a CL < 100 mm possessed elliptical/closed gonopores (Figure 2.5a) indicating that they had not mated, a conclusion consistent with the absence of mating scars in the region of their gonopores (Figure 2.5b) (see also Melville Smith, 1987a; Haefner, 1977). Although the gonopores of female H. acerba did not exhibit such a conspicuous dichotomy in shape as those of C. bicolor, the individuals of this species with a CL < 70 mm had relatively very small gonopores and, unlike many larger individuals, never bore mating scars in the region of their gonopores (Figure 2.6). The first and small group of *H. acerba* was thus considered not to have mated, which is consistent with the ovaries of all such individuals being at stage I (see Table 5.1, Chapter 5 for description of the characteristics of the different stages in ovarian development).



**Figure 2.5.** (a) Abdomen of an immature female *Chaceon bicolor* showing elliptical/closed gonopores and (b) abdomen of a mature female *Chaceon bicolor* showing open and circular gonopores and mating abrasions.



**Figure 2.6.** Abdomen of a mature female *Hypothalassia acerba* showing open and circular gonopores.

# 3.0 THE FISHERIES FOR *HYPOTHALASSIA ACERBA* AND *CHACEON BICOLOR*

#### 3.1 INTRODUCTION

The fishery for the champagne crab *Hypothalassia acerba* is concentrated on the lower west coast of Australia between Jurien in the north at 30° 08'S, 114° 30'E and Mandurah in the south at 32° 29'S and 115° 40'E, and on its south coast between Augusta in the west at 34° 20'S, 115° 10'E and Bremer Bay in the east at 34° 40'S, 119° 30'E (Chapter 1). Under deepsea crab endorsements to fishers by the Western Australian Department of Fisheries, *H. acerba* was only fished commercially on the lower west coast between 1989 and 2000. Although that endorsement extended beyond 2000, the holders of this endorsement switched to targeting the crystal crab *Chaceon bicolor* in that year. The catches of *H. acerba* by fishers with deep-sea crab endorsements have been supplemented to a small degree by those of commercial rock lobster fishers. However, since 2000, the latter fishers have been permitted to retain only six *H. acerba* per vessel per day on the west coast, the champagne crab has been fished commercially since 1991 on the south coast, where this species contributes to a multi-species fishery that also includes the king crab *Pseudocarcinus gigas* and the southern rock lobster *Jasus edwardsii* (Anon., 2004).

*Chaceon bicolor* occurs in water depths of 450 to 1220 m, which is far greater than the depths of 90 to 310 m in which *H. acerba* is typically caught (Melville Smith *et al.*, in press; Chapter 4). The fisheries for *C. bicolor* off the lower west coast and south coast commenced in 1999 and 2002 respectively. Although *C. bicolor* is distributed around Australia and New Zealand, it is only commercially fished in Western Australian waters. However, several other species belonging to the Geryonidae are fished elsewhere in the world (*e.g.* Wigley *et al.*, 1975; Wenner *et al.*, 1987; Erdman and Blake, 1988; Melville Smith, 1988; Defeo *et al.*, 1991;

López Abellán *et al.*, 2002). Although a deep-water crab endorsement entitles fishers to harvest *C. bicolor* as well as *H. acerba*, the former species requires refrigerated water for it to be kept alive and therefore tends to be fished independently of *H. acerba* by fishers who have specialised tanks for holding *C. bicolor*.

The first aim of this chapter is to describe quantitatively the trends exhibited by catch (landed live weight), fishing effort (number of pot lifts) and catch per unit effort (CPUE) for *H. acerba* and *C. bicolor* on the lower west and south coasts since the commencement of the fisheries for these species, and to determine whether these three variables change throughout the year. The second aim was to determine whether the relative abundance and size of both the females and males of *H. acerba* and *C. bicolor* differed among the regions in which they were being fished. The third aim was to tag and release individuals of *H. acerba* in an attempt to obtain data on the movement and growth of this species.

#### 3.2 MATERIALS AND METHODS

#### 3.2.1 Data on commercial catches

The monthly landed catches of each species by commercial fishers on both the lower west and south coasts between 1989 and 2003, together with the number of pot lifts, were extracted by M. Cliff and L. Bellchambers from the Catch and Effort Statistics (CAES) of the Department of Fisheries Western Australia.

#### 3.2.2 Onboard collection of data

#### Hypothalassia acerba

Data for *H. acerba* were collected onboard the boat of a commercial fisher operating in five Department of Fisheries 1° x 1° fishing blocks along the lower west coast and those of three commercial fishers working in five such fishing blocks on the south coast. The first two digits identifying these fishing blocks represent the location of the north-west corner of the block in degrees latitude and the second two numbers represent the last two of the three numbers of the location of the north-west corner of the block in degrees longitude, *e.g.* longitude 114° is given as 14. Fishing blocks 3014, 3114 and 3115 were grouped to represent the northern region and blocks 3214 and 3215 the southern region of the lower west coast, while fishing blocks 3415, 3515 and 3516 were grouped to represent the western region and blocks 3517 and 3419 the eastern region of the south coast. The onboard trips were conducted at two to four month intervals between July 1999 and December 2000 on the lower west coast and less frequently between July 1999 and February 2002 on the south coast (see Chapter 2 for further details). Note that sampling on the south coast occurred less frequently because fishing on this coast was more sporadic and the time spent by fishers at sea was far greater. Whenever possible, a record was kept of the carapace length of each male and female *H. acerba* caught (see Chapter 2) and the latitude and longitude where each line of traps was laid.

#### **Chaceon bicolor**

The onboard collection of the same data for *C. bicolor*, as those recorded for *H. acerba* (see above), was restricted to the west coast as this species was not fished on the south coast during the present study. Catches of *C. bicolor* were grouped into two regions, *i.e.* to the north and south of 28° 50'S (approximately Geraldton), which lies in the middle of the range of latitudes where sampling was undertaken. Onboard recording of *C. bicolor* was undertaken bimonthly between June 2001 and December 2003, with an additional four trips being undertaken in 2000.

#### Mesh selectivity of Chaceon bicolor

During onboard sampling, three recreational rock lobster traps, enclosed with plastic mesh of  $20 \text{ mm} \times 20 \text{ mm}$ , were randomly placed in each of two lines of ~ 100 non-meshed traps of the same type as those used by the commercial fishers who target *C. bicolor* (see Chapter 2.1.6). On each sampling occasion, the carapace lengths of crabs caught in non-meshed and meshed traps were recorded. When the senior author was not present, commercial fishers were asked to record, whenever feasible, the sex and carapace length of crabs in both meshed traps and in an equal number of non-meshed traps. The data recorded by the fishers were later forwarded to Murdoch University for inclusion in our analyses.

Note that the experiments carried out to explore the mesh selectivity of *H. acerba* are reported in Chapter 4 as the data derived from those experiments contributed to the total data set used to determine the depth distribution of this species.

#### 3.2.3 Data Analysis

#### Hypothalassia acerba

Plots of  $log_{10}$  of the standard deviation against  $log_{10}$  of the mean for both the CPUE (catch per pot lift) and the carapace lengths of *H. acerba* recorded on commercial vessels in each of the two regions on each coast showed that neither variable required transformation prior to subjection to analysis of variance (ANOVA) (see Clarke and Gorley (2001) for rationale for this approach).

The catches were subjected to ANOVA, with the number of crabs per pot lift as the dependent variable and sex and region as fixed factors. The carapace lengths were also subjected to ANOVA, with carapace length as the dependent variable, and sex and region as fixed factors. No attempt has been made to separate the data by season because it was not

possible to obtain samples from each region in every season. Scheffé's *a posteriori* test was used to compare the CPUE and mean carapace lengths for each of the four regions.

#### **Chaceon bicolor**

The catch and carapace length data for *C. bicolor* were analysed in the same way as for *H. acerba*. Note however that in this case the catch rates required log transformation.

#### 3.3 RESULTS

#### 3.3.1 Commercial catch and effort data for Hypothalassia acerba

The total landed catch of *H. acerba* in Western Australia rose from less than 1 500 kg in 1989 and 1990 to between 13,050 and 10,850 kg in 1991 to 1996 and then sharply to a peak of 72,975 kg in 1998, before declining progressively to 17,295 kg in 2003 (**Figure 3.1a**). The annual catches peaked on the west coast at 30,900 to 45,630 kg in 1997 to 1999 and on the south coast at 26,070 and 27,345 kg in 1997 and 1998, respectively (**Figure 3.1b**). However, catches on the west coast declined precipitously to essentially zero after 2000, whereas on the south coast they remained at between 10,790 and 21,560 kg between 2000 and 2003 (**Figure 3.1b**).

The total number of pot lifts per annum on both coasts collectively rose progressively from ~5,500 in 1989 and 1990 to 114,850 in 1994 and then declined to 48,220 in 1996, before rising again to reach a maximum in 1998 (**Figure 3.2a**). It subsequently remained at between 44,000 and 61,000 pot lifts in 1999 to 2003. The number of pot lifts per annum on the west and south coasts followed different trends. Thus, on the west coast, they peaked sharply at 73,000 and 94,000 in 1997 and 1998, respectively, and had declined to very low levels by 2001, whereas, on the south coast, they peaked at 90,510 in 1994 and were relatively high in 2001 to 2003 (**Figure 3.2b**).



Figure 3.1. (a) Total landed catch of *Hypothalassia acerba* in south-western Australia and (b) landed catch of this species on the lower west and south coasts of Western Australia separately. Data were extracted from the catch and effort statistics of the Department of Fisheries Western Australia and encompass the years between 1989 and 2003.



**Figure 3.2.** (a) Total number of pot lifts for *Hypothalassia acerba* in south-western Australia and (b) number of pot lifts for this species on the lower west and south coasts of Western Australia separately. Data were extracted from the catch and effort statistics of the Department of Fisheries Western Australia and encompass the years between 1989 and 2003.

The annual catch per unit effort (CPUE) for the whole fishery for *H. acerba*, as expressed by catch (kg) per pot lift, peaked in 1999 at 0.80, after which it declined progressively to 0.25 in 2003 (**Figure 3.3a**). Prior to 1991, the effort on both coasts was either low or zero (**Figure 3.2b**) and thus attention is not focused on the CPUEs for those years. On the west coast, the CPUEs remained at between 0.30 and 0.53 in all years between 1991 and 2000, except in 1995 and 1996 when they were 0.15 and 0.09, respectively, and in 1999 when they rose to a sharp peak at 1.25. The catch rates, and also the effort, on the west coast in 2001 to 2003 were negligible and thus are unlikely to provide reliable measures of relative abundance. The annual CPUE on the south coast rose to a maximum of 0.85 in 1997 and then declined to 0.27 in 2003 (**Figure 3.3b**).

Between 1992 and 1999, when the annual catches of *H. acerba* on the west coast of Australia were greatest, the mean monthly catches on that coast appeared to follow an annual cycle, increasing from low levels of 620 to 1,240 kg in March to June to maximum levels of 1,790 to 3,000 kg in August to January (**Figure 3.4a**). The mean monthly catches on the south coast for the years 1992 to 2003, when catches were substantial on that coast, followed a distinct annual cycle, declining from 2,390 kg in January to a minimum of 410 kg in August, and then rising steeply to 2,260 kg in December (**Figure 3.4b**).

The mean monthly number of pot lifts on the west coast followed a less seasonal cycle than that of total landed catch and only exceeded 4,000 in January and December (**Figures 3.4a, 3.5a**). The mean monthly number of pot lifts on the south coast followed similar trends to those exhibited by the mean monthly catches, declining from 8,085 in January to between 960 and 2,245 in May to October and then rising to 6,445 in December (**Figure 3.5b**).

The mean monthly CPUEs for *H. acerba* on the west coast rose progressively from low levels of < 0.65 kg per pot lift in January to April to reach a maximum of 2.47 kg per pot lift in November, and then declined precipitously to 0.20 kg per pot lift in December (**Figure 3.6a**).


**Figure 3.3.** (a) Total catch per pot lift of *Hypothalassia acerba* in south-western Australia and (b) total catch per pot lift of this species on the lower west and south coasts of Western Australia separately. Data were extracted from the catch and effort statistics of the Department of Fisheries Western Australia and encompass the years between 1989 and 2003.



Figure 3.4. Mean monthly landed weight of *Hypothalassia acerba*  $\pm$  95 % CIs on (a) the lower west coast in the years 1992 - 1999 and (b) the south coast in the years 1992 - 2003.



**Figure 3.5.** Mean monthly number of pot lifts for *Hypothalassia acerba*  $\pm$  95 % CIs on (a) the lower west coast in the years 1992 - 1999 and (b) the south coast in the years 1992 - 2003.



Figure 3.6. Mean monthly CPUEs of *Hypothalassia acerba* + 95 % CIs on (a) the lower west coast between 1992 and 1999 and (b) the south coast between 1992 and 2003.

The mean monthly CPUEs for *H. acerba* on the south coast rose from ~ 0.37 kg per pot lift in January to March to attain a peak of 1.08 kg per pot lift in July (**Figure 3.6b**).

#### 3.3.2 Commercial catch and effort data for Chaceon bicolor

The total annual catch of *C. bicolor* in Western Australia rose progressively from a negligible level in 1997 to reach a maximum of 222,520 kg in 2001 and remained at above 193,280 kg in 2002 and 2003 (**Figure 3.7a**). The vast majority of the catch in each year came from the west coast (**Figure 3.7b**).

The annual number of pot lifts for *C. bicolor* followed a similar trend to that of landed catch, except that they peaked one year earlier, *i.e.* in 2000 rather than 2001 (**Figure 3.8a**). The maximum number of pot lifts per year on the west coast was 245,000.

The CPUE for *C. bicolor* in south western Australia rose progressively from 0.11 kg per pot in 1998 to 1.16 kg per pot in 2002 and 2003 (**Figure 3.9a**). Since the effort was very low on the south coast (**Figure 3.8b**), emphasis was not placed on the catch per unit effort for this coast (**Figure 3.9b**).

The mean monthly catches of *C. bicolor* show no conspicuous seasonal trends, with values ranging from minima of 12,775 to 14,485 kg in April, June and November to maxima of 8,895 to 20,240 kg in January, August, September and October (**Figure 3.10a**). The trends exhibited by the mean monthly number of pot lifts and, to a large extent, the mean monthly CPUE, essentially parallel those of the mean monthly catches (cf **Figures 3.10a,b,c**).

### 3.3.3 Analysis of catches and sizes of crabs derived from onboard sampling

#### Hypothalassia acerba

ANOVA demonstrated that the CPUEs of *H. acerba* differed significantly between sexes and among regions, with the mean square being far higher for sex than region (**Table 3.1**). The sex × region interaction was not significant (**Table 3.1**). The



Figure 3.7. (a) Total landed catch of *Chaceon bicolor* in south-western Australia and (b) landed catch of this species on the lower west and south coasts of Western Australia separately. Data extracted from the catch and effort statistics of the Department of Fisheries Western Australia for the years between 1999 and 2003.



**Figure 3.8.** (a) Total number of pot lifts for *Chaceon bicolor* in south-western Australia and (b) number of pot lifts for this species on the lower west and south coasts of Western Australia separately between 1994 and 2003. Data extracted from the catch and effort statistics of the Department of Fisheries Western Australia



**Figure 3.9.** Total catch per pot lift of *Chaceon bicolor* in south-western Australia and (b) total catch per pot lift of this species on the lower west and south coasts separately for the years between 1994 and 2003. Data extracted from the catch and effort statistics of the Department of Fisheries Western Australia.



**Figure 3.10.** (a) Mean monthy landed weight of *Chaceon bicolor*  $\pm$  95 % CIs on the lower west coast and (b) mean monthy number of pot lifts  $\pm$  95 % CIs on the lower west coast and (c) mean monthy landed weight per pot lift  $\pm$  95 % CIs on the lower west coast. Data were for the years between 2000 and 2003.

mean CPUE of males, *i.e.* ~2.5 per pot, was far greater than that of females, *i.e* ~0.9 (**Figure 3.11a**).

**Table 3.1** Mean squares and significance levels for ANOVA of catch per pot lift (CPUE) of male and female *Hypothalassia acerba* in northern and southern regions off the lower west coast and western and eastern regions off the south coast of Western Australia. Data were recorded onboard commercial vessels between 1999 and 2003. \*\*P<0.01, \*\*\*P<0.001; d.f., degrees of freedom.

	Mean Square	d.f.
Sex (S)	286.4***	1
Region (R)	13.7**	3
S x R	5.4	3
Residual	2.8	990

The mean number of crabs per pot in the four regions ranged from 2.1 in the southern region of the west coast to 1.4 in the western region of the south coast (**Figure 3.11b**). Scheffé's *a posteriori* test showed that the mean CPUE in the southern region of the lower west coast was significantly greater (P < 0.05) than in its northern region.

#### Chaceon bicolor

ANOVA demonstrated that the CPUE for C. bicolor differed significantly between sexes (P < 0.001) and between regions (P < 0.01), with the mean square being greater for sex than for region (**Table 3.2**). The sex × region interaction was not significant (P > 0.05) (**Table 3.2**). The mean CPUE was greater for males than females, *i.e.* 1.4 vs 1.2 per pot (**Figure 3.12a**) and was greater in the northern region than southern region, *i.e.* 1.4 vs 1.3 per pot (**Figure 3.12b**).



**Figure 3.11.** (a) Mean numbers of male and female *Hypothalassia acerba* per pot lift  $\pm$  95% CIs and (b) mean numbers of *Hypothalassia acerba* per pot lift  $\pm$  95% CIs in each region.



Figure 3.12. Mean numbers of *Chaceon bicolor* per pot lift  $\pm$  95% CIs for (a) males and females and (b) the northern and southern regions of the west coast.

**Table 3.2**Mean squares and significance levels for ANOVA of catch perpot (CPUE) of male and female *Chaceon bicolor* in northern and southernregions off the lower west coast. Data were recorded onboard commercialvessels between 1999 and 2003. \*\*P<0.01, \*\*\*P<0.001; d.f., degrees of</td>freedom.

	Mean Square	d.f.
Sex (S)	13.0***	1
Region (R)	2.2**	1
S x R	0.3	1
Residual	0.2	5671

#### 3.3.4 Carapace Lengths

#### Hypothalassia acerba

ANOVA demonstrated that the carapace length of *H. acerba* was influenced significantly by both sex and region (P < 0.001), with the mean square for sex being slightly greater than that for region (**Table 3.3**). The region × sex interaction was not significant.

**Table 3.3.** Mean squares and significance levels for ANOVA of carapace lengths of male and female *Hypothalassia acerba* in northern and southern regions off the lower west coast and western and eastern regions off the south coast of Western Australia. Data were recorded onboard commercial vessels between 1999 and 2003. \*\*\*P<0.001; d.f., degrees of freedom.

	Mean Square	d.f.
Sex (S)	2954.2***	1
Region(R)	2559.7***	3
S x R	104.8	3
Residual	146.2	1713

The mean carapace length of males (100.7 mm) was greater than that of females (95.2 mm) (**Figure 3.13a,b**). The mean carapace length of *H. acerba* ranged from 91 mm in the eastern region of the south coast to 103 mm in the western region of the south coast (**Figure 3.13c**). Scheffé's *a posteriori* test showed that the mean carapace lengths of *H. acerba* differed



Figure 3.13. (a) Mean carapace lengths of male and female Hypothalassia acerba + 95% CIs and (b) Carapace length - frequency distributions and mean carapace lengths for male and female Hypothalassia acerba caught by commercial fishers off the lower west and south coasts of Western Australia and c) mean carapace lengths of Hypothalassia acerba + 95% CIs in each region.

significantly (P < 0.05) between each pair of regions, except for between the southern region of the west coast and the western region of the south coast.

Although the maximum length of males (139 mm) was greater than that of females (120 mm) in the samples examined onboard commercial vessels on the lower west and south coasts, the modal carapace length classes of males and females were both 90-94 mm (**Figure 3.13b**).

#### **Chaceon** bicolor

The carapace lengths of *C. bicolor* differed significantly between the sexes and between the two regions (**Table 3.4**). The mean square was far greater for sex than for region, and the region  $\times$  sex interaction was not significant (**Table 3.4**).

**Table 3.4**Mean squares and significance levels for ANOVA of carapacelengths of male and female *Chaceon bicolor* in northern and southern regionsoff the lower west coast of Western Australia. Data were recorded onboardcommercial vessels between 1999 and 2003. \*\*\*P<0.001; d.f., degrees of</td>freedom.

	Mean Square	d.f.
Sex (S)	72513.2***	1
Region(R)	6298.8***	1
S x R	28.1	1
Residual	323.7	13326

The mean carapace length of the males of *C. bicolor* (119.0 mm) was greater than that of its females (106.3 mm CL) (**Figure 3.14a,b**). Furthermore, males reach a larger maximum size (169 mm) than females (148 mm CL) (**Figure 3.14b**). In contrast to *H. acerba*, however, the modal size class of male *C. bicolor* (115–119 mm) was substantially greater than that of females (100–104 mm) (**Figure 3.14b**). Few crabs were caught with carapace lengths



Figure 3.14. (a) Mean carapace lengths of male and female *Chaceon bicolor* ± 95% CIs, (b) Carapace length - frequency distributions and mean carapace lengths for male and female *Chaceon bicolor* caught by commercial fishers off the lower west coast of Western Australia and (c) mean carapace lengths of *Chaceon bicolor* ± 95% CIs for each region.

< 80 mm. The mean carapace length of *C. bicolor* was greater in the southern than northern region, *i.e.* 113 *vs.* 110 mm (**Figure 3.14c**).

#### 3.3.5 Carapace lengths of Chaceon bicolor in non-meshed and meshed traps

In the case of the males of *C. bicolor*, the mean carapace lengths were greater in the catches obtained in meshed than non-meshed traps in each of the three seasons in which substantial numbers were caught in the non-meshed traps (**Table 3.5**). However in none of these seasons was the difference in size significant. The mean carapace length of females was significantly greater in the catches obtained in the non-meshed than meshed traps in summer (P < 0.05), the only season in which the catches of females in non-meshed traps was substantial (**Table 3.5**). However, even in this latter case, the difference between the carapace lengths in the two trap types was < 8 mm. It is thus concluded that the enclosure of traps in relatively fine mesh did not lead to a conspicuous increase in the capture of small crabs. This point is highlighted by the fact that the percentage of the crabs caught <80 mm CL in the meshed traps were only slightly greater than in the non-meshed traps, *i.e.* 3 vs 1%.

		Males	n	P	Size range (mm)
Summer	Non-mesh	119.6 <u>+</u> 0.7 mm	2115	NS	34 - 169
	Mesh	115.8 <u>+</u> 5.3 mm	63	143	38 - 155
Winter	Non-mesh	117.9 <u>+</u> 1.1 mm	601	NS	59 - 160
	Mesh	112.6 <u>+</u> 8.8 mm	20		82 - 140
Spring	Non-mesh	116.2 <u>+</u> 1.0 mm	785	NS	72 - 153
	Mesh	114.8 <u>+</u> 4.1 mm	45	145	88 - 143
		Females			
Summer	Non-mesh	106.5 <u>+</u> 0.7 mm	782		57 - 141
	Mesh	98.8 <u>+</u> 6.6 mm	47	*	35 - 128

**Table 3.5** Mean carapace lengths  $\pm 95$  % CIs of male and female *Chaceon* bicolor caught in non-meshed vs meshed traps in four seasons between 2001 and 2003. *n*, sample size; NS = not significant at P > 0.05; \* = P < 0.05.

#### 3.3.6 Results from release of tagged Hypothalassia acerba

Twenty eight of the 1622 champagne crabs tagged and released off the lower west coast were recaptured. All of the nine tagged females and nineteen tagged males that were released and recaptured were mature. However, none of the females was ovigerous when it was either tagged, released or recaptured. All but four of the tagged females that were subsequently caught were recaptured within 50 km of their point of release. Two of the recaptured tagged crabs had moved ~ 200 km in a north-westerly direction and a further two had moved over 200 km in a south-westerly direction (**Figure 3.15**). These crabs had thus moved along the contour of the upper shelf break, which runs approximately parallel to the lower west Australian coastline.

None of the 28 recaptured crabs moulted, and this included even one female that had been released back into the environment for over 12 months.

#### 3.4 DISCUSSION

#### 3.4.1 Annual trends in catch statistics

#### Hypothalassia acerba

The data shown in **Figure 3.1** demonstrate very clearly how the overall catches of *Hypothalassia acerba* in Western Australia rose from very low levels in 1989 and 1990 to peak at over 70 000 kg in 1998 and then declined sequentially in subsequent years to under 20 000 kg in 2002 and 2003. Those data also show that the fishery for *H. acerba* on the west coast became essentially non-existent in 2001 to 2003, due to the deep-sea crab fishers on this coast shifting their effort almost exclusively to catching *Chaceon bicolor*.

The trends exhibited by the annual number of pot lifts for *H. acerba* in Western Australia demonstrate that the fishing effort in this state peaked in 1998, when the number of pot lifts was ~ 135 000, and has subsequently remained relatively constant at about half this level from 1999 to 2003. A decline in effort on the west coast in the last five years has been



**Figure 3.15.** Distances in km of the location of recaptured tagged *Hypothalassia acerba* from the point of their release.

compensated for by an increase in effort on the south coast. The decline undergone each year since 1999 by the overall CPUE for Western Australia, as reflected in the trends exhibited by the annual catches per pot lift for the state, indicate that the relative abundance of *H. acerba* has declined during recent years. This suggests that fishing pressure has had an effect on the density of this species.

#### Chaceon bicolor

The data in **Figure 3.7** emphasise that the commercial fishery for *C. bicolor* in Western Australia commenced later than that of *H. acerba*, but that catches rose sharply from negligible values in 1997 and 1998 to peak at 225 000 kg in 2001 and remained close to that level in 2001 to 2003. The rapid increase in the fishery was due to the development of a sudden and focused interest on *C. bicolor*, which led to the construction of vessels that were purpose-built for targeting this species (D. Hand, pers. comm.). The fisheries data also emphasize that the fishery has been based almost exclusively on the west coast. Although the progressive rise in CPUE between 1998 and 2002/3 suggest that the relative abundance of *C. bicolor* increased during this period, that upward trend is almost certainly attributable to a rapid increase in the knowledge by fishers of the best locations and best fishing methods for catching this species.

#### 3.4.2 Monthly trends in catch statistics

#### Hypothalassia acerba

The monthly trends exhibited by the fishing effort for *H. acerba* on the south coast closely paralleled those of catch, with values being least in early winter to mid spring. The relatively low fishing effort in the above period reflects, in part the adverse weather conditions at this time and, also in the case of the months between July and October, the closure of the fisheries for one of the other two species that comprise the multi-species fishery for large crustaceans on the south coast, *i.e.* the southern rock lobster *Jasus edwardsii*. The similarity in the trends exhibited by both catch and effort accounts for the fact that the catch per unit effort did not change markedly throughout the year.

In contrast to the situation with *H. acerba* on the south coast, the monthly trends displayed by effort for this species on the west coast did not parallel those of catch. Indeed, effort was least in September, October and November on this coast, which explains why the catch per unit effort was greatest in these months. Although caution must be exercised in drawing conclusions from these catch statistics for this new fishery, the consistency of the progressive rise in CPUE between summer and the following spring does suggest that the relative abundance, or more likely the catchability of *H. acerba*, is greatest in the spring months.

#### **Chaceon bicolor**

The lack of a conspicuous seasonality in either the catch or fishing effort for *C. bicolor* on the west coast accounts for the CPUE for this species on this coast remaining remarkably constant throughout the year. This, in turn, implies that the relative abundance and catchability of *C. bicolor* on the west coast vary little during the year.

#### 3.4.3 Catches and carapace lengths of Hypothalassia acerba and Chaceon bicolor

The fact that the catches of females were significantly less than those of males in the case of both *H. acerba* and *C. bicolor* parallels the results recorded for the trap catches of other species of deep-sea, crabs such as *Chaceon affinis*, *Chaceon maritae* and *Chaceon notialis*, *Pseudocarcinus gigas*, *Chionoecetes bairdii*, *Chionoecetes tanneri* (Melville Smith, 1988; Defeo *et al.*, 1991; Levings *et al.*, 2001; López Abellán, *et al.*, 2002; Workman and Phillips, 2002; Rosenkranz, 2002). The dominance of males in trap catches has been attributed to such features as male aggression (Bovjberg, 1956) and trap avoidance by ovigerous females (Melville Smith, 1987). Although there were some regional differences in the CPUEs for both *H. acerba* and *C. bicolor*, the differences were not pronounced and the basis for these differences was not clear.

The mean carapace lengths of the males of both *H. acerba* and *C. bicolor* were significantly greater than those of their females, as is the case with several other species of deep-sea crab (Melville Smith, 1988; Levings *et al.*, 1996; López Abellán, *et al.*, 2002). However, the modal carapace length class of males and females of *H. acerba* are the same, *i.e.* 90–94 mm, whereas that of the males of *C. bicolor, i.e.* 115-119 mm, is considerably greater than that of the females, *i.e.* 100–104 mm. The differences between the sizes of the two sexes is thus greater in *C. bicolor* than *H. acerba*.

### 4.0 RELATIVE ABUNDANCES AND SIZE COMPOSITIONS OF *HYPOTHALASSIA ACERBA* ON TWO COASTS AND IN DIFFERENT WATER DEPTHS AND SEASONS

#### 4.1 INTRODUCTION

The champagne crab *Hypothalassia acerba*, together with the western rock lobster *Panulirus cygnus*, the southern rock lobster *Jasus edwardsii*, the king crab *Pseudocarcinus gigas* and the crystal crab *Chaceon bicolor*, are the main large benthic, crustacean predators in water depths greater than 20 m off the south-western Australian coast (George, 1962; Kailola *et al.*, 1993; Levings *et al.*, 1996, 2001; Koh and Ng, 2000; Jones and Morgan, 2002).

The acquisition of appropriate biological data is crucial for managing deep-sea crab species as these grow slowly and are long lived and are thus susceptible to becoming the subject of boom-and-bust fisheries (Hastie, 1995). Prior to the present study, no such data on features such as distribution patterns, size compositions and sex ratios were available for *H. acerba*, or for any other species of the family Eriphiidae except those provided in reports on the king crab *Pseudocarcinus gigas* by Gardner (1998) and Levings *et al.*, (2001).

However, such data are available for a number of deep-sea crab species elsewhere in the world, and particularly for certain species belonging to the Geryonidae and Lithodidae. The distribution of several of these species is known to be related to water depth, temperature and/or substrate type (Melville Smith and Bailey, 1989; Hastie, 1995; Comeau *et al.*, 1998) and the males tend to be more abundant in catches and grow to a larger size than the females (Lux *et al.*, 1982; Melville Smith, 1988; Attrill *et al.*, 1991; López Abellán *et al.*, 2002). Furthermore, body size has been shown to be inversely correlated with water depth in certain deep-sea geryonid species, implying that the individuals of these species are recruited into deeper water and subsequently move shorewards as they increase in size (Beyers and Wilke, 1980; Attrill *et al.*, 1990; Lockhart *et al.*, 1990). In some species, such movements are apparently more pronounced amongst females than males, occur seasonally and/or are related to such factors as temperature and/or reproductive state (Melville Smith, 1987b; Lockhart *et al.*, 1990; Defeo *et al.*, 1991; Hastie, 1995). Moreover, differences between the compositions of the invertebrate fauna living on the shells of males and females of *Chionoecetes opilio* imply that, when not pairing, the two sexes of this species tend to occupy different niches off Newfoundland (Hooper, 1986). There is also evidence that the spatial resources in deep-sea waters are partitioned among two congeneric species of deep-sea crab in the Gulf of Mexico, with *Chaceon quinquedens* being more abundant in shallower waters and *Chaceon fenneri* being more numerous in deeper waters (Lockhart *et al.*, 1990; Hastie, 1995).

During the present study, we collected *H. acerba* during each season from seven water depths located at depth intervals of 55 m between 35 and 310 m on both the lower west and south coasts of Western Australia in order to test the following hypotheses. (1) Hypothalassia acerba will occur predominantly in a range of water depths that differ from those in which the other main large benthic invertebrate carnivores are predominantly found and thus reduce the likelihood of competition for resources amongst those species. (2) Since bottom water temperature declines with increasing water depth and overall is higher at corresponding depths on the lower west coast than south coast, the relative abundance of this species would be likely to peak at a deeper water depth on the lower west than south coast if temperature influences the distribution of *H. acerba*. (3) The distribution by depth will change seasonally in response to changes in bottom water temperature. The study also focuses on determining whether, as with certain representatives of two other families of deep-sea crabs, the following apply on both coasts. The males will be relatively more abundant and attain a greater size than females, the body size will be inversely correlated with water depth and there will be a certain degree of habitat segregation between the sexes. Since the "western rock lobster" traps used in our study caught few small *H. acerba*, as is often the case with the traps employed to catch deep-sea

60

crabs (*e.g.* Lux *et al.*, 1982; López Abellán *et al.*, 2002), the question of whether the enclosure of these traps with a fine mesh would increase the catches of small crabs was also examined.

# 4.2 MATERIALS AND METHODS4.2.1 Sampling Regime

The data used to determine the relative abundance of *Hypothalassia acerba* in different water depths were derived from the depth-stratified sampling regime, which was undertaken once per calendar season on both the lower west and south coasts using traps that were not enclosed in mesh (Chapter 2).

The data used to describe the relationships between carapace length and water depth, coast and season were derived collectively from the sampling described above, but in this case they included the data collected using traps enclosed in small mesh. Refer to chapter 2 for a full description of the sampling regime.

#### 4.2.2 Data Analysis

Plots of  $\log_{10}$  (mean) against  $\log_{10}$  (standard deviation) showed that, prior to subjecting catch rates to Analysis of Variance (ANOVA), they required square root transformation (see Clarke and Gorley (2001) for rationale for this approach). These two values (*mean* and *s.d.*) were calculated from the catches per line of three traps in each water depth on each coast in each season. The transformed values for catch for each sex for each line of traps in each water depth on each coast in each season were subjected to ANOVA, with each factor being fixed. Note that since no crabs were caught at depths of either 35 or 365 m on either coast, these zero values were not employed in the ANOVA. Since the four-way interaction between coast, depth, sex and season was not significant (P > 0.05), this term was removed from the model and the data reanalysed. Results from the ANOVA were back-transformed before plotting. The distribution of crabs with respect to bottom water temperature was explored by fitting a nonlinear regression model that related the catch C(T) by all three lines of traps to the bottom water temperature  $T^{\circ}C$  recorded at the corresponding depth at which the traps were placed. The base model used, *i.e.* 

$$C(T) = a + b \exp\left[-c(T-d)^2\right],$$

was intended to represent catches for the sampling unit that were proportional to a common base level of catch, *a* (where  $a \ge 0$ ), over all temperatures, augmented by a 'normal distribution' of increased catches of crabs centred at a mean temperature of *d*<sup>P</sup>C. The parameters *a*, *b*, *c* and *d* were estimated by minimizing the weighted sum of squared deviations between transformed values of the observed and predicted catches, where each weight was calculated as the inverse of the predicted catch and a penalty function was used to ensure that  $a \ge 0$ . A likelihood ratio test was used to determine whether the above model provided a better description of the distribution of crabs than the simpler model C(T) = a, *i.e.* a distribution of crabs that was independent of water temperature. Likelihood ratio tests were also used to test whether the value of *a* in the base model differed significantly from zero and/or whether there were differences between the mean temperatures of the 'normal distributions' for the curves for the south and west coasts and, on the west coast, between those for the meshed and nonmeshed traps.

The logarithms of the standard deviations of the carapace lengths for each sex on each sampling occasion on both coasts were plotted against the logarithms of the mean values for those carapace lengths. This demonstrated that the carapace lengths did not require transformation prior to subjection to analysis of covariance (ANCOVA). This latter analysis, which employed the General Linear Model (GLM) procedure within SPSS Version 10 (SPSS Inc., Chicago, Illinois, USA) and assumed a normal distribution of observation errors, was

undertaken to ascertain whether there was a significant linear relationship between carapace length and water depth and whether this relationship was influenced by sex, coast and/or season. If the covariate in an ANCOVA is found to be significant, the absolute value of the dependent variable is dependent upon the value of the covariate and values of the dependent variable for the other factors must be adjusted to a standard value of this covariate before they are compared. In the SPSS analysis of the data for *H. acerba*, a water depth of 200 m was specified as the standard depth at which the comparison among other factors was to be made if the covariate, depth, was found to be significant. Likewise, if the other factors (or their interactions) are found in the ANCOVA to be significant, a plot of the values of the dependent variable against the values of the covariate will reflect both the linear trend associated with the covariate and the effects of the other significant factors and interactions. To remove these effects and thereby illustrate more clearly the relationship between the dependent variable and the covariate, the values of the effects of the significant factors and interactions, as calculated in the ANCOVA, were subtracted from the corresponding values of the dependent variable to transform the data to values that would be approximately "equivalent" to those recorded at the reference levels of the factors, *i.e.* females of *H. acerba* on the south coast during spring. The means and 95% confidence intervals of the transformed data were calculated and plotted against the values predicted for the corresponding depths for females on the south coast during spring.

A fully saturated statistical model was fitted to the carapace lengths, with sex, coast and season as factors and depth as a covariate. Higher order interactions that were not significant (P > 0.05) were then successively removed, with the term having the greatest *P*-value being selected as the next term to be eliminated. Next, lower-order interaction terms that were not significant were examined to determine whether they might also be removed to reduce the model further. Such lower-order terms were retained if they contained constituent variables

63

that were included in the remaining significant higher-order interaction terms. The value for the Bayesian Information Criterion (BIC) was calculated for the resulting model as

$$BIC = n \log_e \left(\frac{SS_e}{n}\right) + p \log_e(n)$$
, where  $SS_e$  is the sum of squares of the errors, *n* is the number

of observations, p is the number of parameters that were estimated by the GLM procedure and  $\log_e$  is the natural logarithm (Schwarz, 1978). The highest-order interaction terms in the model were then removed and the BIC recalculated. If the latter value was less than the corresponding value for the preceding model, it was concluded that this latter model was the more appropriate description. In this case, the simpler model was accepted and re-examined to determine whether further non-

significant terms could be removed, using the same procedure as described above.

## 4.3 RESULTS4.3.1 Water temperature

Water temperatures declined with increasing water depth on both the west and south coasts (**Figure 4.1**). However, at each depth for which there are comparable data, the temperatures were appreciably greater on the west than south coast in both summer and spring, but not in autumn or winter. Temperatures in depths  $\leq 255$  m were greater in summer and autumn than in winter and spring on the west coast, whereas they were greater in autumn and winter than in spring and summer on the south coast.

#### 4.3.2 Relative abundance of crabs in different water depths and temperature

ANOVA demonstrated that the catch rates of crabs differed significantly between sexes and among water depths and seasons (**Table 4.1**). However, the mean squares and significance



Figure 4.1Mean bottom water temperatures  $\pm$  95% CIs in the water depths where<br/>Hypothalassia acerba was sampled off the lower west and south coasts<br/>of Western Australia in each season.

levels were far greater for water depth and sex than for season. All of the two-way interactions except sex  $\times$  season and the three-way interactions sex  $\times$  coast  $\times$  water depths and coast  $\times$  depth  $\times$  season were significant (**Table 4.1**).

On both the west and south coasts, the mean catch rates were greater for males than females in those water depths in which the majority of crabs were caught, *i.e.* 200 and 255 m on the west coast and 90, 145 and 200 m on the south coast (**Figure 4.2**). These depth-related differences in the abundances of crabs on the west and south coasts, together with the fact that males were proportionately more abundant on the south coast, explain the sex × coast × depth interaction for catch rates (**Table 4.1**).

**Table 4.1.** Mean squares and significance levels for ANOVA of catch rates of female and male *Hypothalassia acerba* recorded in the seven water depths sampled on the lower west and south coasts of Western Australia during the four seasons of the year. \*P<0.01 \*\*P<0.01 \*\*P<0.001

	Mean squares	d.f.
Sex (S)	16.4***	1
Coast (C)	2.5	1
Depth (D)	23.6***	4
Season (T)	3.1**	3
$S \times C$	7.4**	1
$S \times D$	2.4*	4
$S \times T$	0.2	3
C×D	26.3***	4
C×T	7.1***	3
D × T	7.0***	12
$S \times C \times D$	4.3***	4
$S \times C \times T$	1.2	3
$C \times D \times T$	3.5***	10
$S \times D \times T$	0.8	12
Residual	0.8	54

The catch rates of crabs on the west coast peaked at a water depth of 200 m in summer, autumn and winter and at 255 m in spring (**Figure 4.3**). In contrast, on the south coast, catch rates



**Figure 4.2** Mean numbers of male and female *Hypothalassia acerba*  $\pm$  95% CIs caught in trap lines in different water depths off the lower west (top) and south coasts (bottom) of Western Australia.



Figure 4.3 Mean numbers of *Hypothalassia acerba*  $\pm$  95% CIs caught in trap lines in different water depths off the lower west and south coasts of Western Australia in each season.

peaked at 90 or 145 m in summer, autumn and spring and at 255 m in winter. These differences explain the coast  $\times$  depth  $\times$  season interaction (**Table 4.1**).

Comparison of the fitted model with one that assumed a distribution of catch rates independent of temperature revealed that, on the south coast, catch rates were not significantly influenced by temperature (P>0.05). However, on the west coast, catch rates of champagne crabs were influenced markedly and significantly (P<0.001) by bottom water temperatures (**Figure 4.4**). While the value of the parameter *a* did not differ significantly from zero for either trap type on the latter coast (P>0.05), the mean temperatures of the 'normal distributions' fitted to the numbers of crabs caught using non-meshed and meshed traps, *i.e.* 17.1 and 16.1°C, respectively, differed significantly (P<0.01). Although the temperature distribution of the catches of the non-meshed traps on the south coast were not demonstrated to be significantly influenced by temperature, it should be noted that the mean temperature associated with the 'normal distribution' in the curve fitted to these catches was 16.4°C and did not differ significantly from that for non-meshed traps on the west coast (P>0.05). It is thus concluded that the mean temperature occupied by *H. acerba* of the size range caught by traps on both coasts lies between 16.1 and 17.1°C.

#### 4.3.3 Relationship between carapace length and water depth, coast and season

The carapace lengths of male and female *H. acerba* caught during the study ranged from 50 to 135 mm and from 51 to 113 mm, respectively on the west coast and from 62 to 138 mm and from 69 to 120 mm, respectively, on the south coast (**Figure 4.5**). The size distribution of the females was skewed markedly to the left of males on the south coast but not on the west coast.

As the four-way interaction between the carapace lengths of crabs and sex, coast, depth and season was not significant (P > 0.05) in the ANCOVA to which these data were subjected, the analysis was repeated but excluding this interaction term. Season × depth × sex was the



**Figure 4.4** Distribution of numbers of champagne crabs caught at various bottom water temperatures in non-meshed and meshed traps off the lower west (above) and south (below) coasts of Western Australia.



**Figure 4.5** Frequency distributions for the carapace lengths of male and female *Hypothalassia acerba* caught using non-meshed traps off the lower west and south coasts of Western Australia.

only three-way interaction term that was significant (P < 0.05). However, the value for the \BIC for this model, *i.e.* 5628, was greater than that for the model that excluded this three-way interaction term, *i.e.* 5616. Thus, this three-way interaction term was removed from the ANCOVA, together with non-significant two-way interaction terms.

Carapace length was significantly influenced by water depth and the relationship between these two variables did not exhibit a significant interaction with either sex, coast or season (**Table 4.2**). Thus, on average, the carapace length declined by 4 mm for every 100 m increase in water depth (**Figure 4.6a**). Sex × coast and coast × season were the only significant two-way interaction terms (**Table 4.2**). After adjustment of the data to a standard depth of 200 m, the mean carapace lengths of males and females were 96.6 and 94.6 mm, respectively, on the west coast, and 101.5 and 91.4 mm, respectively, on the south coast (**Figure 4.6b**), the latter difference being significant (P < 0.001). Although the mean carapace length of males on the south coast was significantly greater than that of males on the west coast (P < 0.001), no such significant difference was found between the mean carapace lengths of females on the two coasts (P > 0.05).


Figure 4.6 (a) Mean carapace lengths  $\pm 95\%$  CIs for *Hypothalassia acerba* caught in different water depths off Western Australia. The mean carapace lengths have been calculated after adjusting the data to represent the equivalent lengths that, after adjustment, would have been obtained for females on the south coast during spring. Mean carapace lengths  $\pm$ 95% CIs of (b) male and female *H. acerba* caught off the lower west and south coasts of Western Australia and of (c) both sexes collectively off the lower west and south coasts of Western Australia in each season. The data in Figs 3.5b and 3.5c have been adjusted to a standard depth of 200 m.

**Table 4.2** Mean squares and significance levels for ANCOVA of carapace lengths of female and male *Hypothalassia acerba* caught in the seven water depths sampled on the lower west and south coasts of Western Australia during the four seasons of the year. Depth was treated as a covariate. \*\*\*P<0.001

	d.f.	
Sex (S)	5906.0***	1
Coast (C)	82.0***	1
Depth (D)	2899.7***	1
Season (T)	882.4***	3
$S \times C$	2684.4***	1
$C \times T$	1251.4***	3
Residual	141.0	4

The mean carapace lengths of crabs caught during summer and winter off the south coast were greater than those of crabs caught during these corresponding seasons off the west coast, whereas the reverse pertained in autumn and spring (**Figure 4.6c**). During winter, the mean carapace length of crabs caught off the south coast was significantly greater than that of crabs caught during autumn and spring off that coast (P < 0.001).

#### 4.3.4 Sex ratios

Males dominated the catches obtained from all water depths collectively in each season on both the west and south coasts (**Table 4.3**). However, in each season, the prevalence of males was not as great on the west coast as on the south coast, with the seasonal percentage contributions made by males to the catches of crabs in each season ranging from 64.8 to 76.9% on the west coast compared with 76.0 to 88.8% on the south coast. The overall percentages of males on the west and south coasts were 69.2 and 84.0%, respectively (**Table 4.3**).

	West coast			South coast		
	Males	Females		Males	Females	
	%	%	п	%	%	n
Summer	76.9	23.1	143	87.2	12.8	102
Autumn	64.8	35.2	233	79.0	21.0	100
Winter	70.2	29.8	47	88.8	11.2	160
Spring	68.8	31.2	263	76.0	24.0	75
Overall	69.2	30.8	676	84.0	16.0	437

**Table 4.3.** Percentage of females and males of *Hypothalassia acerba* caught in traps in each season and overall on the lower west coast and south coasts of Western Australia. n = number of crabs.

#### 4.3.5 Results of enclosing traps with mesh

On the two occasions when crabs were caught in non-meshed and meshed traps, the mean carapace length of both male and female crabs obtained using meshed traps was significantly different (P < 0.05) from that of the corresponding sex collected using the same type of trap but without mesh (**Figure 4.7**). However, the differences in the mean carapace lengths of the crabs caught in meshed and non-meshed traps differed by only between 3.9 and 6.9 mm in each of the four comparisons. Furthermore, the number of crabs that were caught with carapace lengths of less than 75 mm still remained low.

#### 4.4 DISCUSSION

#### 4.4.1 Relationship between catch rates and water depth

The sampling regime used in this study was designed to collect *Hypothalassia acerba* from the range of water depths in which preliminary studies had indicated that this species occurred on the lower west and south coasts of Western Australia. Furthermore, particular attention was paid to ensuring that the seven sampling sites were aligned perpendicular to the shoreline and were located sequentially at even depth intervals of 55 m from the shallowest site at 35 m to



**Figure 4.7.** Frequency distributions for the carapace lengths of male and female *Hypothalassia acerba* caught off the lower west and south coasts of Western Australia using meshed and non-meshed traps in May (top) and November (bottom). Sample sizes (n) and mean  $\pm$  95% CIs are given on Figure.

the deepest site at 365 m. The sampling of seven sites (water depths) on both coasts in each season thus produced results that enabled conclusions to be drawn regarding aspects of the spatial and size distributions and sex ratios of *H. acerba*. This species of deep-sea crab was not caught in the shallowest depth and was rarely caught in the deepest waters in any season on either coast, and the catch rates at the intermediate depths rose progressively and then declined sequentially. This demonstrates that our sampling regime essentially encompassed the full range of depths at which at least all but the smaller members of this species occur in the sites sampled. Furthermore, the depths at which *H. acerba* were predominantly caught on both the west and south coasts are entirely consistent with those in which commercial fishers obtained their catches (R. Melville Smith, Department of Fisheries W.A., pers. comm.). The very strong relationship between catch rate and water depth is further illustrated by our finding that, in the case of the most abundant sex, *i.e.* males, the catches peaked sharply at 200 m on the west coast and at 145 m on the south coast. Furthermore, the catches of males at 200 and 255 m collectively on the west coast and at 90, 145 and 200 m collectively on the south coast contributed 90 and 92%, respectively, to the total catches of this sex on these two coasts. Although the females of *H. acerba* were less abundant than their males on both the west and south coasts, our data still clearly demonstrate that the trends exhibited by the abundance of females with water depth were the same as those for males on both coasts. Thus, the catches of females likewise peaked at 200 m on the west coast and at 145 m on the south coast. The presence of the same pattern of distribution of the two sexes of *H. acerba* in all seasons contrasts with the situation with certain other deep-sea crabs species in which, when unpaired, the sexes tend to occupy different water depths (Beyers and Wilkie, 1980; Melville Smith, 1987; Lockhart et al., 1990; Defeo et al., 1991). For example, the females of Geryon maritae, Chaceon fenneri, Chaceon notialis, and Chionoecetes opilio were relatively more

prevalent than males in shallower waters (Melville Smith, 1987, 1988; Lockhart *et al.*, 1990; Defeo *et al.*, 1991).

Comparisons of our results with those for other species of deep-sea crabs suggest that *H. acerba* is restricted to a particularly narrow range of water depths for such species. For example, *Chaceon notialis* was caught in water depths of 300 to 900 m in the Argentinian–Uruguayan fishing zone and, in summer, the catch per unit of effort (CPUE) was still increasing at the upper end of this range in water depths (Defeo *et al.*, 1991). Although the catch rates of *Chaceon affinis* peaked at 800-900 m water depth, this species was caught in water depths of 500 to 1200 m and, from the data presented, presumably well beyond that latter depth (López Abellán *et al.*, 2002). *Geryon quinquedens* and *Geryon maritae* are also found in a wide range of water depths, *i.e.* from 275 to 1000 m off southern New England and from 270 to 924 m off Namibia (Lux *et al.*, 1982; Melville Smith and Bailey, 1989).

Since *H. acerba* appears to be largely restricted to water depths within the relatively narrow ranges of 145 to 310 m on the west coast and of 90 to 200 m on the south coast, it is relevant that other major large benthic crustaceans in this region are most abundant in water depths to either side of those in which *H. acerba* occurs. Thus, on the west coast, the western rock lobster *P. cygnis* is most common in depths up to 90 m (George, 1962), while off the south coast the southern rock lobster *J. edwardsii* is also most frequently caught in waters less than 90 m (Department of Fisheries, Western Australia, Catch and Effort Statistics, unpublished data). The crystal crab *Chaceon bicolor* is most abundant in depths exceeding 600 m on the west coast and 500 m on the south coast (K. Smith and R. Melville Smith, unpublished data), where our unpublished sediment analyses showed that the silt/clay fraction in the substrate was far greater than in the shallower waters where the champagne crab is found, *i.e.* 32 vs 5%. Furthermore, on the south coast, *P. gigas* also occurs mainly in water depths that, from Levings *et al.*, (1996, 2001) and Gardner (1996), lie at the upper end of the

range of that in which *H. acerba* lives. The distribution patterns of these decapod species thus provide evidence of spatial partitioning among these major components of the benthic communities on the two coasts, thereby parallelling the type of partial partitioning by depth exhibited by *Chaceon fenneri* and *Chaceon quinquedens* in the Gulf of Mexico (Lockhart *et al.*, 1990).

#### 4.4.2 Relationship between catch rates and water temperature

Since the density of *H. acerba* is related to water depth on both the west and south coasts and water temperature decreases with increasing water depth, it seems highly likely that the relative abundance of this species will also be influenced by water temperature. Such a conclusion is consistent with the fact that, while the overall catches of *H. acerba* peaked at a shallower water depth on the south coast (145 m) than lower west coast (200 m), those peaks in abundance corresponded to a similar water temperature, *i.e.* ~16.1-17.1°C. Thus, by typically occurring in a shallower water depth on the south coast. It is thus not surprising that the distribution of catch rate of *H. acerba* by depth on the two coasts was most similar in winter when the water temperatures at the corresponding depths on the two coasts were most similar.

Although *H. acerba* was caught in water temperatures ranging from 11.9 to 21.9°C on the west coast and from 11.2 to 20.6°C on the south coast, over 85% of the individuals of this species were caught in temperatures of 15.8 to 18.2°C on the west coast and over 95% were collected at temperatures between 13.1 and 18.7°C on the south coast. Furthermore, the water temperature at which the catches of *H. acerba* peak lies between 16.1 and 17.1°C on both coasts, which is ~6 to 10°C higher than those recorded for five *Chaceon* species, which live in deeper water (see Hastie, 1995).

#### 4.4.3 Trap selectivity

Our failure to catch small *H. acerba* in traps parallels the situation recorded for several other species of decapod (Melville Smith, 1986; Miller, 1990). Furthermore, the mean size of crabs caught by the rectangular traps was only slightly reduced when these traps were enclosed in aviary mesh. This may reflect either trap shyness by small *H. acerba* or their exclusion from traps through the aggression of larger individuals or possibly their occupation of different water depths. The greater prevalence of males than females in our catches, and those of commercial fishers operating over a wide geographical area (K. Smith, unpublished data) may reflect females being less vulnerable to traps, as is the case in some other species of decapod (see Miller, 1990). Furthermore, the fact that only seven ovigerous *H. acerba* were caught during the present study, which is again consistent with the results for some other decapod species (Miller, 1990) and presumably reflects a tendency for the berried females of this species not to enter traps, also helps account for the predominance of males in our catches of *H. acerba*.

This chapter has been accepted for publication as follows: Smith, K.D, Hall, N.G. and Potter, I.C. (2004). Relative abundances and size compositions of male and female champagne crabs, *Hypothalassia acerba* (Brachyura : Eriphiidae), on two coasts and in different water depths and seasons. *Mar. Freshwat. Res.* (in press).

# 5.0 COMPARISONS BETWEEN THE REPRODUCTIVE BIOLOGY OF FEMALES OF TWO SPECIES OF DEEP SEA CRABS WHICH LIVE IN DIFFERENT WATER DEPTHS

#### 5.1 INTRODUCTION

Three species of deep-sea crab are fished commercially in Western Australian waters, namely the champagne crab *Hypothalasia acerba* Koh and Ng, the crystal crab *Chaceon bicolor* Manning and Holthuis and the king crab *Pseudocarcinus gigas* 

Lamarck (Anon., 2002). On the basis of catches obtained by commercial fishers, the distribution of *H. acerba* ranges from a latitude at least as far north as about 27°S on the west coast, southwards and then eastwards along the south coast to a longitude of at least 129°E (Anon., 2002). Although *C. bicolor* occurs in the same geographical region as *H. acerba*, its distribution in Western Australia extends further north to a latitude of about 22°S (MelvilleSmith *et al.*, in press). In contrast, the distribution of *P. gigas* is essentially restricted to the south coast in Western Australia and extends eastwards across the full length of Australia, including Tasmania, and then northwards on the east coast to about 31°S (Kailola *et al.*, 1993, Gardner, 1998).

Recent depth-stratified sampling off the south-western Australian coast has demonstrated that *H. acerba* occurs predominantly in waters where temperatures lie between 13 and 20°C and the depths range from 200 to 255 m on the lower west coast and from 90 to 200 m on the south coast (Smith *et al.*, 2004a). Temperatures recorded during commercial fishing for *C. bicolor* show that, on the lower west coast of Australia, this species typically occurs in waters of 4 to 6.5°C (K. Smith unpublished data) and at water depths of 450 to 1220 m (Melville Smith *et al.*, in press) and is thus located further offshore than *H. acerba*. *Pseudocarcinus gigas* is found in waters of 11 to 17°C and at depths of 120 to 340 m (Levings *et al.*, 1996).

81

The commercial fisheries for *H. acerba* and *C. bicolor* in Western Australia started to develop during the late 1990s (Anon., 2002). However, in contrast to the situation with *P. gigas* (see Levings *et al.*, 1996, 2001; Gardner, 1997, Gardner and Williams, 2002), no biological studies have been conducted on the spawning period and location and fecundity of either of these two species.

The stages in development of the ovaries of several species of deep-sea crabs belonging to the Geryonidae, *e.g. Chaceon quinquedens* (Smith), *Chaceon maritae* (Manning & Holthuis) and *Chaceon affinis* (Milne Edwards & Bouvier), have been described on the basis of their macroscopic appearance and, in some cases, also on the stages of maturation of their oocytes as determined from histological sections (Haefner, 1977, Melville Smith, 1987a, Fernández-Vergas *et al.*, 2000, López Abellán *et al.*, 2002). No such combined macroscopical and histological studies have been carried out on any deep-sea species of the Eriiphidae, the family to which *H. acerba* belongs. Although some deep-sea crab species, such as *Chaceon fenneri* (Manning & Holthuis), have an annual reproductive cycle (see Hastie, 1995), no clear seasonal fluctuations in reproductive patterns were detected with *C. maritae* (Melville Smith, 1987a) and *C. quinquedens* (Wigley *et al.*, 1975, Haefner, 1978). It was proposed by Melville Smith (1987a) that the lack of a distinct reproductive cycle in *C. maritae* was attributable to the absence, in the deep waters in which this species is found, of changes in day length and temperature and other potential environmental reproductive triggers.

The aims of the present study were as follows. (1) Describe, on the basis of their macroscopic appearance, the sequential stages in the development of the ovaries of *H. acerba* and use histological sections to determine the stages of maturation of the oocytes in ovaries at each of those stages. Attention was paid to elucidating whether the criteria developed for categorizing the ovarian stages of *H. acerba* were the same as those we developed concomitantly for *C. bicolor* and those described for other species of deep-sea crab. (2) Use the

82

trends exhibited throughout the year by the monthly prevalences of female crabs with ovaries at different stages in development and of females bearing either eggs or egg remnants, together with those of the mean monthly gonad weights of females, to determine the timing of reproduction in these two species. Particular emphasis is placed on testing the hypothesis that, as *H. acerba* lives in far shallower waters than *C. bicolor*, and thus lives in an environment which undergoes a greater degree of seasonal change and particularly in water temperature, it is far more likely to reproduce seasonally. (3) Determine the fecundities of *H. acerba* and *C. bicolor* and their relationships to the body size and body weight of these species. (4) Establish whether aspects of the reproductive biology of *H. acerba* on the lower west and southern coasts of Western Australia exhibit similar patterns. Since preliminary analyses indicated that *H. acerba* invests less in gonadal development on the south than west coasts, we examined whether the relative ovarian weights and diameters of the mature oocytes of females on the two coasts differed.

## 5.2 MATERIALS AND METHODS

#### 5.2.1 Collection of samples

The reader is referred to Chapter 2 for details of the depth-related sampling regime and the onboard sampling which together yielded the *H. acerba* and *C. bicolor* used for the present study.

#### 5.2.2 Laboratory processing

The reader is referred to Chapter 2 for description of measurements made on *H. acerba* and *C. bicolor* and the criteria used to designate female crabs as either immature or mature.

The ovary of each female was removed and allocated to a numerical developmental stage on the basis of macroscopic criteria (see Results). The stages in development of the testes

and vas deferentia were recorded during another study aimed at determining the size at maturity of the males of these two species (Chapter 7).

The ovary of each female of *H. acerba* and *C. bicolor* was weighed to the nearest 0.1 g. The mean monthly ovarian weights ( $\pm$  95% CIs) at standardised carapace lengths were determined for *H. acerba* on the lower west and south coasts and for *C. bicolor* on the former coast using analysis of covariance (ANCOVA) of the natural logarithm of the gonad weight as the dependant variable, month as a fixed factor and the natural logarithm of the carapace length as a covariate. Note that the standardised carapace length was calculated separately for the lower west and south coast assemblages of *H. acerba* and the lower west coast assemblage of C. bicolor. These constant carapace lengths were a default value calculated by the ANCOVA. ANCOVA was also used in the above manner to calculate the mean gonad weight ( $\pm 95\%$  CIs) for *H. acerba* with ovaries containing previtellogenic, early vitellogenic and late vitellogenic oocytes on the lower west and south coasts. The standardised carapace length was calculated separately for each of these three developmental stages, but was common for a given stage on the two coasts. Data for the small individuals that were considered not to have mated were not used when calculating the monthly prevalences of females with ovaries at different stages in development, the mean monthly gonad weights for females at a standardized carapace length, and the mean gonad weights for west and south coast H. acerba with ovaries at different stages in development (see Figures 5.2-5.5 in Results).

The method of Melville Smith (1987a) was used for determining the batch fecundity of *H. acerba* and *C. bicolor*. Thus, alternate pleopods were removed from the abdomen of each ovigerous female and soaked for 1 to 4 h in 0.5 M NaOH to facilitate the separation of eggs from the pleopod setae. All of the eggs removed from each crab were pooled, oven dried at 70°C for 24 h and then weighed to the nearest 0.001 g. Subsamples of ca. 0.1 g of eggs (= ca. 300 eggs) were then weighed and counted, and the average weight of a single egg in each

subsample was calculated. When estimates of the weight of a single egg in three subsamples differed by less than 5%, the mean of those weight estimates was then used, in conjunction with twice the weight of all eggs removed, to estimate the fecundity of the crab.

The right posterior side of the ovary was removed from up to 20 females of both species in each season, ensuring that these subsamples covered the full size range of those species in the total catch in each season. The ovarian material was fixed in 4% glutaraldehyde in a 0.025 M phosphate buffer solution (pH 7.0), dehydrated in a series of increasing concentrations of ethanols, embedded in paraffin wax, sectioned transversely at 6 µm and stained with Mallory's trichrome. The ovarian material was fixed in glutaraldehyde since it was not fixed well by traditional fixatives used for light microscopy such as Bouin's and 10% formalin. The smallest and largest diameters of oocytes of *H. acerba* and *C. bicolor* at different developmental stages sectioned through their nuclei were measured to the nearest 5 µm. The smallest and largest diameters of each oocyte were then averaged.

In addition to the above laboratory processing of crabs, the senior author also recorded, when on board fishing vessels, the total number of mature female crabs caught and the percentage of both ovigerous females and of females with egg remnants attached to their pleopods. Note that, due to inclement weather, the commercial fisher with whom we mainly worked did not operate in September in the deep waters off the lower west coast of Australia where *C. bicolor* is found and thus there are no data on the above variables for that month.

#### 5.3 RESULTS

#### 5.3.1 Ovarian maturation stages

On the basis of their macroscopic appearance, the ovaries of the females of *Hypothalassia acerba* and *Chaceon bicolor* each followed essentially the same pattern of development (**Table 5.1**). Thus, for both species, six sequential stages were recognised (**Table 5.1**; Figure 5.1). As

**Table 5.1.** Macroscopic and microscopic characteristics of sequential stages in ovarian development of *Hypothalassia acerba* and *Chaceon bicolor*. Range in oocyte diameters at each stage of development are provided. Scheme has been adapted from that of Haefner (1977).

Ovarian stage	Macroscopic description	Histological description
<b>Stage I</b> Virgin/immature resting	Ovaries small but generally visible macroscopically, thin and 'H' shaped. (Shape is maintained for all latter stages except stage IV). Oocytes not visible through ovarian wall. Colour varies from pale grey to ivory.	Ovaries contain oogonia and small previtellogenic oocytes (chromatin nucelolar and perinucleolar oocytes). Connective tissue and oocytes stain blue with Mallory's trichrome. For <i>H. acerba</i> , oogonia 5-10 µm and small previtellogenic oocytes 10-60 µm. For <i>C. bicolor</i> , oogonia 5- 10 µm and small previtellogenic oocytes 10-70 µm.
Stage II Developing	Ovaries always visible macroscopically, larger than stage I and slightly dorso-ventrally compressed. Ovaries beginning to displace hepatopancreas. Oocytes not visible through ovarian wall. Colour varies from ivory to white.	Ovaries contain all oocyte stages present in stage I together with cortical alveolar oocytes. Cortical alveolar oocytes 60-200 $\mu$ m in <i>H. acerba</i> , 70-210 $\mu$ m in <i>C. bicolor</i> . Cortical alveolar oocytes stain blue with Mallory's trichrome, except for their lipid droplets and nucleolei, which stain dark red.
<b>Stage III</b> Maturing	Ovaries larger than at stage II and outer membrane has wrinkled appearance. Oocytes not visible through ovarian wall. Ovaries are pale pink in <i>H. acerba</i> while range from very pale yellow to yellowish orange in <i>C. bicolor</i> .	Large numbers of early yolk granule oocytes present. Yolk granules in peripheral region of oocyte cytoplasm. Yolk granules stain orange with Mallory's trichrome and are larger than lipid droplets. For <i>H. acerba</i> , yolk granule oocytes are $160 - 270 \mu\text{m}$ and for <i>C. bicolor</i> are 140-390 $\mu\text{m}$ .
Stage IV Mature	Ovaries larger than at stage III and consequently are no longer 'H- shaped'. Ovarian wall more folded. Oocytes visible through ovarian wall. Ovaries pink to red in <i>H. acerba</i> and brown to purple in <i>C. bicolor</i> .	Large numbers of late yolk granule oocytes. Yolk granules distributed throughout cytoplasm. For <i>H. acerba</i> yolk granule oocytes are 200- 480 µm and for <i>C. bicolor</i> 350-560 µm.
Stage V Recently spent	Ovaries similar in appearance to stage II, but more flaccid and yellowish-grey in colour.	Ovary contains mainly connective tissue. Some yolk granule oocytes present undergoing atresia.
Stage VI Spent-recovering	Same as stage V.	Ovaries comprise mainly connective tissue and numerous small previtellogenic oocytes.



**Figure 5.1.** Histological sections of ovaries of *H. acerba* at (a) stage I, (b) stage II, (c) stage III, (d) stage IV, (e) stage V (recently-spent) and (f) stage VI spent-recovering. a, atretic oocyte; ca, cortical alveolar oocyte; ct, connective tissue; eyg, early yolk granule; o, oogonia; lyg, late yolk granule; pv, pre vitellogenic oocyte. Scale bar =  $200 \mu m$ .

they progress from stages I to IV, the ovaries increase progressively in size and change colour (**Figures 5.1a-d**). However, the colours of the ovaries of the two species at stages III and IV differ. In both species, ovaries at stages V and VI are flaccid and could not be separated macroscopically.

At a histological level, cortical alveolar oocytes first appeared in stage II ovaries of *H*. *acerba* (**Figures 5.1a,b**), and the presence of early and late yolk granule oocytes characterised stage III (**Figure 5.1c**) and IV ovaries (**Figure 5.1d**), respectively. Ovaries of recently-spent females (stage V) of *H. acerba* contained some atretic oocytes (**Figure 5.1e**), while those of spent-recovering (stage VI) females possessed no such oocytes but did contain numerous early previtellogenic ooctytes (**Figure 1f**). The histological characteristics of the ovaries of *C. bicolor* are essentially the same as those shown for *H. acerba* in **Figures 5.1a-f**. However, the mean diameters of the mature oocytes, as measured in stage IV ovaries, are significantly greater (P < 0.001) for *C. bicolor* than *H. acerba*.

Comparisons of the macroscopic and microscopic characteristics of our ovarian stages for *H. acerba* and *C. bicolor* with those of Haefner (1977) demonstrate that the ovaries of the former two species undergo the same pattern of development as *Chaceon quinquedens*. However, the maximum diameter of the oocytes in our stage I (70 µm) is far less than that recorded by Haefner (1977) for his first stage for *C. quinquedens* (172 µm) and yolk granule oocytes are already present in his stage II whereas they do not appear until stage III of our categorization for *H. acerba* and *C. bicolor*. In essence, Haefner (1977) did not recognise the presence of the early stage we have termed virgin/immature resting. However, that stage was recognised by de Lestang *et al.* (2003) in their description of ovarian development in *Portunus pelagicus* (Linnaeus) in inshore waters in Western Australia. Furthermore, unlike the study of Haefner (1977), our histological staging criteria distinguish recently-spent ovaries from spentrecovering ovaries, which is useful for obtaining an idea of how recently ovulation had occurred.

#### 5.3.2 Monthly trends in reproductive indices

On the lower west coast, the prevalence of female *H. acerba* with ovaries containing late vitellogenic oocytes was greatest in September to February, during which months their contributions ranged from 58 to 76%, and was least in March to June when their monthly contributions ranged from 11 to 50% (**Figure 5.2a**). The prevalence of female crabs with ovaries containing only previtellogenic oocytes was less than 12% in each month from September to February and in April and only exceeded 30% in March and August. Small numbers of spent females were collected between January and March and in May. The mean monthly gonad weight of *H. acerba*, standardized to a constant carapace length of 94.7 mm, rose progressively from 11.5 g in July to between 25.2 and 28.6 g in October to December and then declined sequentially to between 4.9 and 8.3 g in March to May (**Figure 5.2b**). The prevalence of female *H. acerba* bearing egg remnants declined sequentially from 12% in July to zero or close to zero in October to January before rising progressively to 21% in May. Small numbers of ovigerous females were caught in February, March and May (**Figure 5.2c**) and a commercial fisher provided six ovigerous females and two females with egg remnants which he had caught in January 2000.

On the south coast, the prevalence of female *H. acerba* containing late vitellogenic oocytes and the mean monthly gonad weights, standardized to a common carapace length of 95.2 mm, peaked in September (**Figures 5.3a-b**). The prevalence of *H. acerba* with ovaries containing late vitellogenic oocytes never exceeded 47% in any month other than September and the prevalence of ovaries with previtellogenic oocytes exceeded 50% in five months

89



Month

Figure 5.2. Monthly data for female *Hypothalassia acerba* on lower west coast of Australia. (a) Frequency of occurrence of crabs at different stages of development, (b) mean gonad weights  $\pm$  95 CIs standardised to a constant carapace length of 94.7 mm, as selected by ANCOVA and (c) frequency of occurrence of crabs with eggs and egg remnants. In this Fig. and Figs 3 and 4, the sample sizes for each month in (b) are the same as those given in (a).



Figure 5.3. Monthly data for female *Hypothalassia acerba* on the south coast of Australia. (a) Frequency of occurrence of crabs at different stages of development and (b) mean gonad weights  $\pm$  95 CIs standardised to a constant carapace length of 95.2 mm, as selected by ANCOVA.

(**Figure 5.3a**). Samples from the south coast yielded no ovigerous females and only two females bearing egg remnants, both of which were caught in August.

The prevalences of females of *C. bicolor* with ovaries containing late vitellogenic oocytes were greater in July to December than in January to April and in June (**Figure 5.4a**). The mean monthly gonad weights, standardized to a constant carapace length of 110.0 mm, peaked at 42.2 g in September and 43.1 g in December before declining to 11.6 g in February, increasing to 37.1 g in May and then declining to 8.8 g in June (**Figure 5.4b**). Ovigerous females were caught in each of the 11 calendar months in which samples were obtained and females bearing egg remnants were found in six of those months (**Figure 5.4c**).

The mean gonad weight  $\pm 95\%$  CIs, standardized to a constant carapace length, did not differ significantly (P > 0.05) between the assemblages on the lower west and south coasts when the females contained ovaries with either previtellogenic or early vitellogenic eggs (**Figure 5.5**). In contrast, the mean gonad weight  $\pm 95\%$  CIs for females of *H. acerba* with ovaries containing late vitellogenic oocytes was significantly greater (P < 0.001) for the assemblage on the lower west coast,  $26.4 \pm 1.71$  g, than that on the south coast,  $16.0 \pm 1.85$  g (**Figure 5.5**).

The relationships between gonad weight (*W*) and carapace length (*CL*) of west coast and south coast females of *H. acerba* with ovaries containing late vitellogenic oocytes (**Figure 5.6**) are described by the following equations.

West coast:  $\ln(W) = 2.561 \ln(CL) - 8.130$ ; P < 0.001,  $R^2 = 0.240$ , n = 228South coast:  $\ln(W) = 6.430 \ln(CL) - 26.311$ ; P < 0.001,  $R^2 = 0.442$ , n = 34

Although none of the females of *H. acerba* from the south coast, which had CLs < 90 mm, contained ovaries with late vitellogenic oocytes, many females of *H. acerba* from the lower west coast with CLs of 75 to 90 mm possessed ovaries with oocytes at this advanced stage. The use of ANCOVA demonstrated that, in the case of *H. acerba* for a standardized



## Month

Figure 5.4. Monthly data for female *Chaecon acerba* on the lower west coast of Australia. (a) Frequency of occurrence of crabs at different stages of development and (b) mean gonad weights  $\pm$  95% CIs standardised to a constant carapace length of 110.0 mm, as selected by ANCOVA and (c) frequency of occurrence of crabs with eggs and egg remnants.



Figure 5.5. Mean gonad weights  $\pm 95\%$  CIs for female *Hypothalassia acerba* caught off the lower west versus south coasts and containing ovaries predominantly previtellogenic, early vitellogenic and late oocytes. The gonad weights of each of the ovarian categories on both coasts were standardised to constant carapace lengths as selected by ANCOVA.



**Figure 5.6.** Relationship between the weights of ovaries containing late vitellogenic oocytes and carapace lengths for *Hypothalassia acerba* from both the lower west and south coasts of Western Australia.

selected carapace length, the mean gonad weight of crabs was significantly greater on the lower west coast than south coast (P < 0.001).

To enable predictions of *W* from *CL* for *H. acerba*, the regression equations relating these two variables are provided below,

West coast:  $\ln(W) = 2.809 \ln(CL) - 6.400$ ; P < 0.001,  $R^2 = 0.946$ , n = 581

South coast:  $\ln(W) = 2.768 \ln(CL) - 6.128$ ; P < 0.001,  $R^2 = 0.893$ , n = 185

The mean lengths ( $\pm 95\%$  CIs) of females from the lower west ( $93.8 \pm 2.02$  mm) and south coasts ( $93.0 \pm 1.39$  mm) were not significantly different (P > 0.05) when the individuals contained previtellogenic oocytes, whereas the corresponding values of  $94.7 \pm 0.91$ mm and  $99.1 \pm 1.37$  mm for females with late vitellogenic oocytes on those two coasts were significantly different (P < 0.001). These differences are reflected in the fact that the distributions of the carapace lengths of female *H. acerba* on the south coast were located more to the right than those on the lower west coast in the case of females with late vitellogenic oocytes, whereas this was not so with females with previtellogenic oocytes (**Figures 5.7a,b**).

The mean diameters ( $\pm$  95% CIs) of late vitellogenic oocytes of *H. acerba* from the lower west coast (357.7  $\pm$  10.79 µm, *n* = 342) and south coast (370.3  $\pm$  8.16 µm, *n* = 134) were not significantly different (*P* > 0.05).

The fecundities of *H. acerba* ranged from 202 295 to 488 758 for crabs with CLs of 90 mm and 102 mm, respectively. The minimum and maximum fecundities for *C. bicolor* were 15 592 and 288 512 respectively, for crabs with CLs of 98 and 133 mm respectively. The mean fecundities ( $\pm$  95% CIs) of *H. acerba*, *i.e.* 356 210 ( $\pm$  64 297) and *C. bicolor*, *i.e.* 192 070 ( $\pm$  33 640), were significantly different (*P* < 0.001). Although the mean body weights ( $\pm$  95% CIs) of the ovigerous females of *H. acerba* used for the above fecundity estimates, *i.e.* 587.4 ( $\pm$  96.5 g) were substantially less than that of *C. bicolor*, *i.e.* 651.9 ( $\pm$  61.3 g), those means were not significantly different (*P* > 0.05). The relationships between fecundity (*F*) and the carapace



**Figure 5.7.** Carapace length-frequency distributions for female *Hypothalassia acerba* with ovaries containing (a) previtellogenic and (b) late vitellogenic oocytes off the lower west and south coasts of Western Australia.

length (*CL*) and wet weight (*W*) of the two species are shown in **Figures 5.8a-d** and are described by the following regression equations:

Hypothalassia acerba

 $\ln F = 3.44 \ln(CL) + 1.235, P < 0.05, R^2 = 0.486, n = 10$ 

 $F = 0.005W + 0.608, P < 0.05, R^2 = 0.569, n = 10$ 

Chaceon bicolor

 $\ln F = 2.957 \ln(CL) + 1.686, P < 0.05, R^2 = 0.127, n = 40$ 

 $F = 0.003W + 0.1572, P < 0.05, R^2 = 0.260, n = 40$ 

#### 5.4 DISCUSSION

#### 5.4.1 Comparisons between Hypothalassia acerba on the west and south coasts

The trends exhibited by the three reproductive variables recorded for the females of *Hypothalassia acerba* on the lower west coast of Australia were consistent with each other and clearly imply that reproduction in this deep-sea species is highly seasonal. Thus, the increase in the prevalence of females with ovaries containing late vitellogenic oocytes between July and October and subsequent levelling off to December was paralleled by a progressive rise and then levelling off in the mean monthly relative gonad weights. Likewise, the subsequent decline in the prevalence of females with this category of advanced ovary between December and March was accompanied by a pronounced decline in the mean monthly relative gonad weights. Furthermore, spent females were found in the relatively short period of January to March, which yielded most of the ovigerous females obtained during this study. It is also highly relevant that the prevalence of female *H. acerba* bearing egg remnants rose progressively from zero or very low levels in October to January to peak in May and subsequently declined sequentially to zero between July and October. The consistent trends exhibited by the monthly data for the three reproductive variables strongly suggest that oviposition occurs mainly in January to March.



**Figure 5.8.** Relationships between fecundity and carapace length and body weight for *Hypothalassia acerba* and *Chaceon bicolor*.

The patterns exhibited by the reproductive variables for female *H. acerba* on the south coast of Western Australia differed markedly from those displayed by this species on the lower west coast. Thus, on the south coast, the prevalence of females with late vitellogenic oocytes and the mean monthly relative gonad weights both peaked in September rather than December and the overall prevalence of females with ovaries containing previtellogenic oocytes was far higher. Moreover, no ovigerous female crabs and only two females with egg remnants were collected on the south coast. This strongly suggests that very little reproduction occurs on the south coast.

The possibility that the lack of capture of ovigerous females of *H. acerba* on the south coast is due to females moving inshore prior to oviposition, as occurs, for example, with *Chaceon affinis, Chaceon quinquedens, Chaceon maritae* and *Chionoecetes opilio* (Fabricus) (Haefner, 1978; Hastie, 1995; Melville Smith, 1987a, Comeau *et al.*, 1998, López Abellan *et al.*, 2002) appears unlikely. This conclusion is based on the results of our stratified sampling in sequential water depth intervals from 35 m to 365 m, which showed that the females and males of *H. acerba* were both very largely concentrated in a relatively narrow range of water depths of 200 to 255 m off the lower west coast and of 90 to 200 m off the south coast, and that no crabs were caught in water depths of 35 m (Smith *et al.*, 2004a). However, it is relevant that, although the catches of *H. acerba* on both coasts contained a greater number of males than females, that difference was substantially greater on the south coast (Smith *et al.*, 2004a). The above data on *H. acerba* suggest that, at some stage prior to oviposition, the females of *H. acerba* on the south coast typically migrate westwards and then northwards on the lower west coast where they extrude their eggs.

The conclusion that the females of large and maturing *H. acerba* undergo considerable migratory movements is consistent with the observation that such movements are undertaken, in particular, by mature females of some other species of deep-sea crabs, *e.g. Chaceon fenneri*,

100

C. quinquedens and C. maritae (Melville Smith, 1987b, Lockhart et al., 1990). Some species of lobster also move substantial distances, including the western rock lobster Panulirus cygnus George along the west coast of Australia (Phillips, 1983; Melville Smith and Cheng, 2002). In decapods, these types of movements are often related to reproduction and enable those species to return to their spawning locations as mature adults after they have been transported as larvae or juveniles away from the areas of spawning (e.g. Bennett and Brown, 1983; Moore and Macfarlane, 1984; Stewart and Kennelly, 1988; Groeneveld and Branch, 2002). In the context of possible larval movement in south-western Australia, it is relevant that the important Leeuwin current flows southwards along the lower west coast of Australia and then eastwards along the south coast and extends to a depth of 350 m (Pearce and Griffiths, 1991). It is thus proposed that the larvae of H. acerba are planktonic, as has been shown by Gardner (1998) to be the case with those of the confamilial *Pseudocarcinus gigas*, and would thus be likely to be transported by this major current from the areas on the lower west coast where spawning occurs. The likelihood of such transport would be enhanced by the fact that the larvae of *H. acerba* on the west coast are released in autumn, when the Leeuwin current is strongest (Phillips *et al.*, 1991).

#### 5.4.2 Comparisons between Hypothalassia acerba and Chaceon bicolor

The trends exhibited by the reproductive variables for *Chaceon bicolor* were far less seasonal than those displayed by *H. acerba*. Thus, although the prevalence of females of *C. bicolor* with ovaries containing late vitellogenic oocytes was high in October to December, it was also relatively high in August and May and, while the mean monthly relative gonad weight declined between December and February, it remained high from July to December. More importantly, however, *C. bicolor* with spent/spent-recovering ovaries were caught in eight months and in each season of the year, thereby indicating that oviposition occurs in most months of the year.

This conclusion is consistent with the fact that ovigerous females of *C. bicolor* were caught in each of the eleven months for which samples were obtained for deriving this variable and that, unlike the situation with *H. acerba*, the prevalence of such females did not exhibit a clear pattern of change during the year. The finding that, unlike the situation with *H. acerba*, and also *P. gigas*, which lives in similar water depths (Levings *et al.*, 1996), *C. bicolor* clearly does not reproduce within a restricted part of the year is consistent with *C. bicolor* occupying far greater depths and thus living in an environment that undergoes little seasonal change. The lack of conspicuous seasonality in the reproduction of *C. bicolor* parallels that found by Melville Smith (1987a) for *Chaceon maritae*, which likewise lives in deep water. It is also consistent with the fact that the physiological processes of the majority of organisms living in deep waters do not exhibit seasonality (Tyler, 1988).

#### 5.4.3 Investment in gonadal development by Hypothalassia acerba

It was particularly noteworthy that the mean relative gonad weights of females with late vitellogenic oocytes were far greater for the assemblage on the west coast than for that on the south coast, especially as this did not apply to females with oocytes in earlier stages of development. This strongly indicates that the females of *H. acerba* on the west coast invest relatively less energy in ovarian development than those on the south coast. Furthermore, the lack of a significant difference between the size of the late vitellogenic oocytes of females on the south and lower west coasts implies that any differences in energy allocation to ovarian development on the two coasts is not reflected in differences in an allocation to individual oocytes. From the above data and comparisons, it appears reasonable also to assume that the number of eggs per batch produced by females of *H. acerba* that originated on the south coast are likely to be less than those of the individuals that spent at least most of their life on the west coast.

It is also evident from the data shown in **Figure 5.6** that the females of *H. acerba* on the south coast do not start producing mature (late vitellogenic) oocytes until they have reached a substantially larger size than those on the west coast. This implies that, with respect to size, the attainment of maturity by the females of *H. acerba* is delayed on the south coast compared with the west coast (69 mm CL).

In summary, our results provide good circumstantial evidence that conditions are less conducive for gonadal development and reproduction by *H. acerba* on the south than lower west coast of Western Australia. Furthermore, they also suggest that maturing females of *H. acerba* migrate from the south coast, which is at the southern and cool end of the distribution of this species, to the west coast for spawning. Reproduction on the lower west coast was shown to occur seasonally in *H. acerba*, but throughout at least much of the year with *C. bicolor*, which occurs in far deeper water and thus where environmental conditions remain more constant during the year. The seasonality of reproduction by *H. acerba* is accompanied by producing, at any one time, a greater number of eggs per given body weight than does *C. bicolor*.

This chapter has been accepted for publication as follows: Smith, K.D., Potter I.C. and Hesp, S.A. (2005). Reproductive biology of females of two species of commercially fished crabs, *Chaceon bicolor* and *Hypothalassia acerba*, in waters off south-western Australia. J. Shellfish Res. (in press).

# 6.0 POTENTIAL BIAS IN ESTIMATES OF THE SIZE OF MATURITY OF CRABS DERIVED FROM TRAP SAMPLES

#### 6.1 INTRODUCTION

The age at which fish and crustaceans are first caught should be at least as great as the age at first maturity if recruitment overfishing is to be avoided (Caddy and Mahon, 1995). Therefore, the size at which the individuals of a decapod species typically reach the onset of sexual maturity (SOM) is frequently used by fisheries managers as the basis for assigning a minimum legal size for the retention of that species (Watson, 1970). The criteria used to designate whether a female decapod is mature vary among species, and often reflect differences in the secondary sexual characteristics of those species. For example, because the abdomen of female portunid crabs becomes relatively wider and more loosely attached at the pubertal moult, this characteristic provides a sound criterion for assessing whether a female of this family has reached maturity (*e.g.* Van Engel, 1958; Somerton, 1981; de Lestang *et al.*, 2003a), whereas the shape of the gonopores (Melville Smith, 1987), or the presence of eggs under the abdomen (Levings *et al.*, 2001), have been employed for this purpose with the females of some deep-sea species of crabs.

The size at which 50% of the females of the stock of a crab species reach the onset of sexual maturity (SOM<sub>50</sub>) is usually estimated by subjecting the proportions of mature females in sequential size classes to logistic regression analysis (*e.g.* Fisher, 1999; Muino *et al.*, 1999; de Lestang *et al.*, 2003a). The validity of this approach depends on both the immature and mature individuals of the stock in question having been sampled in an unbiased manner. As many crab fisheries employ only baited traps, scientists have often used the data obtained from the resulting catches to estimate the SOM<sub>50</sub> (Brown and Powell, 1972; Melville Smith, 1987; Abbe, 2002; Gardner and Williams, 2002). However, traps yield samples that are biased in

terms of both the size and sex composition of the population (Williams and Hill, 1982; Zhou and Shirley, 1997; Jury *et al.*, 2001).

Trap selectivity for decapod species typically occurs as a result of the larger individuals being more aggressive, thereby restricting the likelihood of smaller animals entering the trap (*e.g.* Bovbjerg, 1956; Rodhouse, 1984). In addition, the greater number of males than females typically found in the trap catches of decapods (Carroll and Winn, 1989; Potter and de Lestang, 2000) almost certainly reflects the greater aggression of males. Moreover, behavioural studies strongly indicate that, in the case of the portunid *Callinectes sapidus*, females are more likely to enter traps containing males once they have become mature (Jivoff and Hines, 1998). Hence, trap catches of the females of this species will almost certainly contain a disproportionately greater number of mature than immature females. Therefore, subjection of the proportions of trap-caught mature females of *C. sapidus* in successive size classes to logistic regression analysis would presumably underestimate the SOM. Similar biases in the estimates of SOM would be expected to result from analyses of trap catches of other crab species if there is likewise a greater likelihood of the capture of the mature females of those species than their immature females of the same size.

The commercial fishery for the blue swimmer crab (*Portunus pelagicus*) in Western Australia is the largest for this portunid in Australia, a total catch of 673 t being taken in 1999– 2000, yielding a wholesale value of approximately \$A3 million (Anon., 2002). The SOM of female *P. pelagicus* in Shark Bay, the location of the largest fishery for this portunid in Western Australia, has been estimated using data from samples obtained by baited traps, seinenetting and otter trawling collectively (de Lestang *et al.*, 2003a). However, because of trap selectivity, the samples of *P. pelagicus* collected by baited traps in Shark Bay may have contained an atypically high proportion of mature females and, consequently, a logistic regression analysis using data from trap catches may have underestimated the SOM of the females there. In contrast, the size composition of samples of *P. pelagicus* collected by seinenetting and otter trawling would be far more representative of the population in Shark Bay.

The champagne crab (*Hypothalassia acerba*) and the crystal crab (*Chaceon bicolor*) are fished commercially using baited traps in water depths of about 150-360 m and 600-800 m, respectively, off the west and south coasts of Western Australia (Smith *et al.*, in press; unpublished data). Although estimates of the SOMs can be derived from the trap catches of the females of these populations, no data are available from catches obtained using other fishing gear that could be used to determine whether vulnerability to capture by traps was greater among the mature females of these species.

This study aimed to determine whether the SOM<sub>50</sub> estimated for female *P. pelagicus* in Shark Bay was significantly lower when data derived from trapping were used in preference to data from seine-netting and otter trawling collectively. If this was the case, it would demonstrate that, as with *C. sapidus*, the mature female *P. pelagicus* are more vulnerable to capture by traps than the immature females of the same size. Consequently, a SOM<sub>50</sub> derived from trap catches will be an underestimate for species such as *P. pelagicus* and *C. sapidus*. However, the only method used for catching *H. acerba* and *C. bicolor* in Western Australia is trapping. Although it is recognized that the SOM<sub>50</sub>s derived from these trap catches may represent underestimates, we derived such values so that fisheries managers have data that, with discretion, can be used for determining, for example, the minimum legal size for capture of these species.

# 6.2 MATERIAL AND METHODS6.2.1 Sampling and measurement of crabs

The data for *P. pelagicus* used in this study were extracted from the database constructed for the samples of this species collected in Shark Bay (26°S, 113.5°E), at bimonthly intervals between July 1998 and May 2000 (de Lestang *et al.*, 2003a, b). Samples were collected with a

seine-net 21.5 m long with a bunt made of 3 mm mesh, a small otter trawl with a net containing a codend constructed of 25 mm mesh, and a series of crab traps consisting of either 12 or 76 mm mesh (for further details see Potter and de Lestang, 2000). The crab traps, which were the same as those used by most professional fishers in Western Australia, were ca. 630 mm high and 1000 mm in diameter. They were joined together in four lines of four, separated by a distance of 15 m. The traps were baited with fish, usually *Sardinops neopilchardus*, and set for ca. 24 h. The carapace width (CW) of each female *P. pelagicus*, *i.e.* the distance between the tips of the two lateral spines of the carapace, was measured to the nearest 1 mm.

The *H. acerba* and *C. bicolor* used in this study were derived from the depth-stratified sampling regime and from commercial catches as described in Chapter 2. That chapter also describes the measurements made on these two species of crab.

As the morphology and tightness of the abdominal flap of the immature and mature females of both *H. acerba* and *C. bicolor* do not differ markedly, abdominal flap characteristics could not be used to determine the maturity status of the females of those deep-sea crabs. Moreover, as the morphology of the gonopores of immature and mature females of *H. acerba* does not differ markedly, which contrasts with the situation in some other species of deep-sea crab (Melville Smith, 1987), this structure could not be used to determine the maturity status of the champagne crab. Therefore, female *H. acerba* were classified as having reached maturity when their ovaries were relatively large and pale yellow to pink (and shown by histology to contain oogonia and primary oocytes). As the gonopores of female *C. bicolor* are elliptical and compressed in immature individuals, and circular and open in mature individuals, it was possible to use this criterion to determine whether an individual crab was immature or mature.

#### 6.2.2 Data analysis

The SOM<sub>50</sub>s of female *P. pelagicus* taken by trap, and by seining and trawling collectively, were estimated using logistic regression analysis of the proportions mature at each CW. The probability that the  $j^{th}$  crab is mature (P<sub>j</sub>) is

 $P_{j} = 1/ \{1 + \exp[-\ln(19)(CW_{j} - SOM_{50})/(SOM_{95} - SOM_{50})]\}$ 

where  $CW_j$  is the CW of that crab,  $SOM_{50}$  and  $SOM_{95}$  the CWs at which 50 and 95% of the assemblage reach maturity, and ln is the natural logarithm of the specified value. This equation represents a simple reparameterization of the logistic equation, which employs the parameters  $SOM_{50}$  and  $SOM_{95}$  that may be more readily interpreted than the traditional parameters of this equation. The  $SOM_{50}$  and  $SOM_{95}$  were determined through use of a computer program that maximizes log-likelihood, and employs AD Model Builder (Fournier, 2000). The confidence limits of these parameters and of the estimated percentages of mature crabs at different CWs were determined from likelihood profile distributions calculated using AD Model Builder. Similarly, estimates of the  $SOM_{50}$ s were derived for trap-caught females of both *H. acerba* and *C. bicolor*. Note that, in the case of each of these latter species, CW is replaced by CL in the above equations.

A likelihood-ratio test (see Cerrato, 1990) was used to determine whether the  $SOM_{50}s$  of *P. pelagicus* estimated from samples caught by trap, and by seining and trawling collectively, were significantly different.

### 6.3 RESULTS

#### 6.3.1 Size composition

The 272 female *P. pelagicus* caught in traps in Shark Bay, of which only 2.9% were immature, ranged from 74 to 178 mm CW (**Figure 6.1a**). In contrast, the size range of the 238 female *P. pelagicus* caught by seining and trawling collectively, of which as many as 57% were immature, ranged from 12 to 160 mm CW (**Figure 6.1a**). The mean CWs of mature females


**Figure 6.1** Size compositions of immature and mature female *Portunus pelagicus* caught in (a) traps, and (b) seine-nets and otter trawls collectively, in Shark Bay, Western Australia.

collected by trapping (127.5 mm), and by seining and trawling collectively (125.5 mm), were not significantly different (P > 0.05).

The CLs of the 598 female *H. acerba* and the 2981 female *C. bicolor* caught by traps ranged from 50 to 114 mm and from 34 to 148 mm, respectively (**Figure 6.2**). The size distributions of both of these species were essentially unimodal, peaking at ~ 95 and ~ 110 mm CL, respectively. The number of immature individuals in the samples of female *H. acerba* and *C. bicolor* were 20 and 186, respectively, thus contributing only 3.3 and 6.2% to the total catch of the females of these two species (**Figure 6.2**).

#### 6.3.2 Size at maturity

A logistic regression analysis of the proportions of mature female *P. pelagicus* in sequential CWs in trap samples yielded a SOM<sub>50</sub> of 86.1 mm, with lower and upper 95% confidence limits of 59.7 and 97.2 mm, respectively (**Figure 6.3a**). A far greater SOM<sub>50</sub>, *i.e.* 101.1 mm, and a far narrower 95% confidence interval, *i.e.* 96.1-105.4 mm, were obtained when subjecting the proportions of mature crabs in seine and trawl samples to the same analysis (**Figure 6.3b**). The above two SOM<sub>50</sub>s were significantly different (*P* < 0.001). As the mean CW of ovigerous female *P. pelagicus* in trap catches did not differ significantly (*P* > 0.05) from that recorded in seine and trawl catches collectively, the size frequency data for these methods were pooled. The CWs of ovigerous females caught in Shark Bay ranged from 104 to 157 mm, with a mean and 95% confidence limits of 131.3  $\pm$  3.4 mm (**Figure 6.3c**).

Logistic regression analyses of the proportions of mature females of *H. acerba* in sequential CLs in samples collected by traps yielded a SOM<sub>50</sub> of 69.7 mm (**Figure 6.4a**). The CLs of ovigerous females and females with egg remnants collectively ranged from 75 to 114 mm, producing a mean and 95% confidence limits of 96.6  $\pm$  3.0 mm (**Figure 6.4b**). Subjection of the proportions of the mature females of *C. bicolor* in trap catches to logistic regression



**Figure 6.2** Size compositions of immature and mature females of (a) *Hypothalassia acerba*, and (b) *Chaceon bicolor* in samples collected by traps on the lower west coast of Australia.



**Figure 6.3** Logistic regression analysis of the proportions of mature female *Portunus pelagicus* caught in (a) traps, and (b) seine and otter trawl nets collectively. (c) Size composition of ovigerous female *P. pelagicus* caught using all three sampling methods. The estimates of the SOM<sub>50</sub>s and 95% confidence limits are denoted by the arrows and dashed lines, respectively. Values for the SOM<sub>50</sub>s and 95% confidence limits are also presented.



**Figure 6.4** (a) Logistic regression analysis of the proportions of mature female *Hypothalassia acerba* caught in traps. (b) Size composition of ovigerous female *H. acerba*. The estimate of the SOM<sub>50</sub> and 95% confidence limits are denoted by the arrows and dashed lines, respectively. Values for the SOM<sub>50</sub> and 95% confidence limits are also presented.

analysis yielded a SOM<sub>50</sub> of 90.5 mm (**Figure 6.5a**). The CLs of ovigerous females and females with egg remnants collectively ranged from 91 to 140 mm, with a mean and 95% confidence limits of  $108.2 \pm 1.0$  mm (**Figure 6.5b**).

#### 6.4 DISCUSSION

The proportion of small female *P. pelagicus* obtained by trapping in Shark Bay is far less than that obtained by seine-netting or otter trawling in the same environment. Indeed, the percentage contribution made by female *P. pelagicus* with a CW < 100 mm to the total catch of females was only 3.3% in trap catches, whereas it exceeded 57.6% in the combined catches obtained by seine-netting and otter trawling. The proportion of large female crabs that entered traps was also high in the case of the two species of deep-sea crabs. Therefore, the proportions of female *H. acerba* < 70 mm CL and of *C. bicolor* < 90 mm CL among all females of each of those species were only 3.3 and 4.4%, respectively. It should be noted that the numbers of small deep-sea crabs caught were low, even when the traps were enclosed in a smaller mesh or plastic covering and had their entrances narrowed.

The size composition data for *P. pelagicus* for Shark Bay, and also for the Leschenault Estuary much farther south (Potter and de Lestang, 2000, unpublished data), demonstrate that, in contrast to mature females, few immature females of this species enter traps. This finding is consistent with the implications of the results of behavioural studies on *Callinectes sapidus* by Jivoff and Hines (1998), which indicated that an immature female of this portunid would be more likely to avoid any male(s) of this species present in traps than would a mature female. Furthermore, male American lobster (*Homarus americanus*) do not allow the immature females of the same species to occupy their shelter (Salmon, 1983). If the proportion of mature females in any size category is greater in trap samples than in the environment, the SOM<sub>50</sub> derived from such samples would underestimate this parameter as a result of the sample being biased towards those mature individuals. This conclusion is strongly supported by the fact that



Figure 6.5 (a) Logistic regression analysis of the proportions of mature female *Chaceon bicolor* caught in traps. (b) Size composition of ovigerous females of *C. bicolor*. The estimate of the SOM<sub>50</sub> and 95% confidence limits are denoted by the arrows and dashed lines, respectively. Values for the SOM<sub>50</sub> and 95% confidence limits are also presented.

the estimated SOM<sub>50</sub> of female *P. pelagicus*, based on data from trapping (86.1 mm CW), was significantly less than that of 101.1 mm CW estimated with the data derived from seine-netting and otter trawling, which would have caught a far more representative sample of the population. The CW of the smallest ovigerous female *P. pelagicus*, 104.0 mm, was substantially greater than the SOM<sub>50</sub> of 86.1 mm, estimated from the proportions of mature females in trap catches, which, as argued above, is an underestimate.

As with *P. pelagicus*, the estimates of SOM<sub>50</sub> for *H. acerba* and *C. bicolor*, 69.7 and 90.5 mm, respectively, calculated using the proportions of mature females in trap catches, were lower than the sizes of the smallest ovigerous females. Catches of those species obtained using active fishing methods are not available, so precluding investigation of the possibility that the vulnerability of the females of these species might be greater for mature than for immature crabs. Until this hypothesis can be tested for each species, the values of the SOM<sub>50</sub>s for *H. acerba* and *C. bicolor* calculated from trap catches should be considered lower "bounds" of the true SOM<sub>50</sub>s rather than accurate estimates of these crucial reproductive parameters.

This chapter has been accepted for publication as follows: Smith, K.D., Hall, N.G., de Lestang, S. and Potter, I.C. (2004). Potential bias in estimates of the size of maturity of crabs derived from trap samples. *ICES* (in press).

## 7.0 DO THE CHELAE OF THE MALES OF THREE CRAB SPECIES UNDERGO ALLOMETRIC CHANGES THAT CAN BE USED TO DETERMINE MORPHOMETRIC MATURITY?

#### 7.1 INTRODUCTION

The size at which the individuals in fished stocks become mature is one of the most important parameters used by managers to develop plans for ensuring the sustainability of those stocks. Indeed, concerns that the reduced number of males found in crab fisheries, that exploit only or predominantly males, might influence the sustainability of the stocks of certain species have highlighted the need for information on the sizes at which the males of those species attain maturity (Conan and Comeau, 1986; Ennis *et al.*, 1988; Gardner and Williams, 2002).

Studies aimed at determining the size at maturity of male crabs have used data on whether individual crabs have become functionally, physiologically and/or morphometrically mature (*e.g.* Goshima *et al.*, 2000; Conan *et al.*, 2001; Comeau and Conan, 1992; Gardner and Williams, 2002; de Lestang *et al.*, 2003a). Evidence of functional maturity is derived from observations that copulation has occurred (Goshima *et al.*, 2000; Gardner and Williams, 2002), whereas demonstration that males are physiologically mature is provided by their production of spermatophores (Warner, 1977; Melville Smith, 1987). The attainment of morphometric maturity by a male crab is identified by determining whether certain body parts, *e.g.* chelae, undergo a conspicuous change at the pubertal moult (Somerton, 1981; Comeau and Conan, 1992; Gardner and Williams, 2002).

A male crab must become both functionally and physiologically mature before it can reproduce and the ability to mate may depend on the crab having become morphometrically mature. In an aquarium study involving males of *Chionoecetes opilio* with carapace widths (CWs) in excess of the size at physiological maturity and mature females, precopulatory pairing behaviour was found to be initiated only by the larger of the males that had become morphometrically mature, *i.e.* > ~95 mm CW (Conan and Comeau, 1986). It was thus concluded that, although physiological maturity is attained by the males of *C. opilio* before they reach a CW of ~60 mm, functional maturity is not attained until individuals have moulted to become morphometrically mature and attained a CW of ~95 mm. However, Paul (1992) has queried why physiologically mature males would produce sperm if they were not to be used immediately. Subsequently, Sainte-Marie *et al.*, (1995) found that males of *C. opilio* from the Gulf of Saint Lawrence became physiologically mature at a CW of only 38.5 mm and that, in non-competitive aquarium studies, males that were physiologically mature but morphometrically immature were functionally mature, *i.e.* mated successfully.

The changes undergone in the allometry of a body part of a male crab may not be sufficiently well defined to be able to be used with confidence for estimating the size at which morphometric maturity is attained (Clayton, 1990; Goshima *et al.*, 2000). For example, in some decapod species, the changes in the "level of allometry" *sensu* Hartnoll (1978) are gradual and the logarithms of the measurements appear to follow a curvilinear trend. Consequently, the straight lines that are fitted to the presumed juvenile and adult data may be artefactual and cannot thus provide a reliable means for distinguishing between pre- and postpubertal moult animals (Conan *et al.*, 2001). Indeed, Somerton (1980) has pointed out that it is essential to test whether a difference between the patterns of growth of that body part in juveniles and adults can be detected.

Somerton (1980) fitted separate, overlapping lines, rather than a single straight line, to the logarithms of the dimensions of various body parts of juvenile and adult crabs and used a *F*-test to determine whether the difference between the sums of squares of those two models was statistically significant. However, the objective function calculated by Somerton (1980), using his iterative approach to fit separate overlapping lines, relies on the allocation of intermediate points to one or other of the straight lines, based on the value of the dependent

118

variable and the predicted values of the lines at that body size. It was thus inappropriate to employ statistical tests, such as the *F*-test, to compare his models of differing complexity using the value of this objective function and that obtained from a traditional regression model. However, several earlier studies on the maturity of male crabs, including our own on *P. pelagicus* (de Lestang *et al.*, 2003a), have used an overlapping-lines model without validating that it was appropriate.

In Western Australia, the blue swimmer crab *Portunus pelagicus* is fished in estuaries and protected coastal waters, whereas the champagne crab Hypothalassia acerba and the crystal crab Chaceon bicolor are fished predominantly in water depths of 90 to 310 m and 450 to 1220 m, respectively (de Lestang et al., 2003b; Smith et al., 2004a; Melville Smith et al., in press). The main aim of the current study was to determine statistically whether the allometry of the chelae of the males of *P. pelagicus*, *H. acerba* and *C. bicolor* underwent a change that could be assumed to reflect a morphological transition from an immature to a mature state. The Akaike and Bayesian Information Criteria were thus used to explore which of a linear, quadratic, broken stick or overlapping-lines model was the most appropriate for representing the relationship between the natural logarithms of the length of the dorsal propodus and body size of the male crabs of each species (Akaike, 1973; Schwartz, 1978). A linear model would imply that allometry does not change as body size increases, whereas a quadratic model would indicate that the pattern of growth changes continuously with increasing body size and the broken stick or overlapping-lines models would imply that there is a distinct change in allometry during growth. If there was found to be a change in allometry, the resulting estimate of the body size at which this change occurred was compared with the size at which the animal becomes physiologically mature. This required the determination of the size at which physiological maturity is attained by the males of *H. acerba* and *C. bicolor* to provide data,

119

which would complement that already derived for this variable for *P. pelagicus* (de Lestang *et al.*, 2003a).

### 7.2 MATERIAL AND METHODS

#### 7.2.1 Collection of crabs

Male *P. pelagicus*, with CWs ranging from 21-154 mm, were collected from Cockburn Sound (32°12'S, 115°48'E) at monthly intervals between February 1997 and March 2000. The crabs were caught using a 21.5 m seine net with a bunt made of 3 mm mesh and a small otter trawl net with a cod end constructed of 25 mm mesh (see de Lestang *et al.*, 2003a).

The *H. acerba* and *C. bicolor* used for this study were collected during the depthstratified sampling regime and sampling onboard commercial fishing boats (for details, see Chapter 2).

#### 7.2.2 Measurements

The carapace width of each male *P. pelagicus*, *i.e.* distance between the tips of the two lateral spines of the carapace, and the carapace length of each male *H. acerba* and *C. bicolor, i.e.* distance across the gastric region from the midpoint between the bases of the two anterior medial spines and the posterior margin of the carapace, were each measured to the nearest 1 mm. Carapace length rather than carapace width is typically measured in deep-sea crabs (*e.g.* Levings *et al.*, 1996; Gardner, 1997; Goshima *et al.*, 2000) as it overcomes the problems of using the distance between the two lateral spines of the carapace, which are particularly prone to wear. For all three species, the length of the dorsal propodus from the proximal edge of the depression below the upper articulation knob to the proximal edge of the depression at the articulation with the moveable finger of the molariform cheliped of each male was measured to the nearest 0.1 mm using vernier calipers. The choice of chela propodus length to provide data aimed at determining whether or not changes in allometry occur was based, in part, on the

demonstration by Gardner and Williams (2002) that this was the most suitable of several variables for assessing morphometric changes in the giant crab *Pseudocarcinus gigas*. The latter species belongs to the same family (Eriphiidae) as *H. acerba*.

#### 7.2.3 Relationship between propodus length and body size

The natural logarithms of the lengths of the dorsal propodus of the males of each species were plotted against the natural logarithms of body size, *i.e.* carapace length or width. In each case, the trend exhibited by the points with increasing body size appeared continuous and without marked disjunction. This suggests that the data for each species might be adequately described by a linear or quadratic regression or a broken-stick regression, *i.e.* a piecewise linear function that is continuous at the point of intersection, or a model comprising two overlapping lines in combination with a logistic function. The logistic function determines the probability that an observation is drawn from the distribution of points around either the first or the second of these overlapping lines. The straight line implies that there is no change in the level of allometry, whereas the quadratic model represents a continuous and smooth change in the level of allometry as body size increases. The broken stick and overlapping-lines models represent two distinct and constant levels of allometry, reflecting the different patterns of growth of morphometrically immature and mature crabs. For the broken stick model, the change occurs when the animals attain the length corresponding to the point at which the two lines intersect. For the model using overlapping lines, the change is gradual and its rate is determined by the parameters of the logistic curve that determine the proportion of the mixture of points associated with each of the overlapping lines. The linear and quadratic models are written as  $\ln DP = a + b \ln S$  and  $\ln DP = a + b \ln S + c(\ln S)^2$ , respectively, and the broken-stick model as

121

$$\ln DP = \begin{cases} d + e \ln S & \text{if } \ln S \le X \\ d + eX + f(\ln S - X) & \text{otherwise} \end{cases}$$

The overlapping-lines model is

$$\ln DP = (1 - P_{S})(a' + b' \ln S) + P_{S}(c' + d' \ln S),$$

where

$$P_{S} = \{1 + \exp[-\ln(19)(S - S_{50})/(S_{95} - S_{50})]\}^{-1}.$$

In these equations, DP = length of the dorsal propodus, S = body size (*i.e.* CL or CW), *a*, *b* (and *c*) = parameters of the linear (and quadratic) model(s), *d* and *e* = intercept and slope, respectively, of the first line, and *f* = slope of the second line of the broken-stick model, X = value of ln *S* at the point of intersection of the two straight lines that form the broken stick, *a'* and *b'* = intercept and slope, respectively, of the first of the overlapping lines, *c'* and *d'* = intercept and slope of the second of the overlapping lines, *P<sub>s</sub>* = probability that a crab with body size *S* is associated with the second of the overlapping lines, *S<sub>50</sub>* and *S<sub>95</sub>* = body sizes at which 50 and 95%, respectively, of crabs are associated with the second of the overlapping lines, *a* and ln = natural logarithm of the associated variable.

It was assumed that observation errors  $\varepsilon$  were independent, identically distributed random variates drawn from a normal distribution, *i.e.* where  $\varepsilon \sim N(0, \sigma^2)$ . Ignoring constants, the log-likelihood was calculated as  $\lambda = -\frac{n}{2} \ln\left(\frac{SS}{n}\right)$ , where *n* is the number of observations and *SS* is the sum of squared deviations of the observed value of  $\ln DP$  from the value predicted using each model. The models were fitted to the data for each species using Solver in Microsoft Excel, and maximising the log-likelihood for each model to estimate the two parameters of the linear model, *a* and *b*, the three parameters of the quadratic model, *a*, *b* and *c*, the four parameters of the broken-stick model, *d*, *e*, *f* and *X* and the six parameters of the overlapping-lines model, *a'*, *b'*, *c'*, *d'*, *S*<sub>50</sub> and *S*<sub>95</sub>. 95% confidence intervals for each parameter in each model were estimated as the 2.5 and 97.5 percentiles of 200 estimates of each of the parameters that were obtained by random resampling, with replacement, of the data for each species and then refitting each model to the resulting set of data. When fitting the overlapping-lines model, a penalty function was used to ensure that, for crabs with a body size less than a specified minimum or greater than a specified maximum, the values of  $S_{50}$  and  $S_{95}$ were such that  $(1 - P_s) > 0.99$  or  $P_s > 0.99$ , respectively. This emulates the approach used by Somerton (1980) whereby crabs with sizes below or above some intermediate size range were assumed to be immature or mature, respectively. For *P. pelagicus*, the values of these specified minimum and maximum body sizes were set at 60 and 110 mm CW, respectively, as these were the values used by de Lestang *et al.*, (2003a). For *H. acerba* and *C. bicolor*, the values specifying this intermediate size range, *i.e.* 67 to 110 mm CL and 81 to 134 mm CL, respectively, were selected from plots of the data for each species and appeared to bracket the range likely to contain the point of inflection of the lines, if such a point exists.

As the four models are not nested, they cannot be compared using the likelihood-ratio test to determine the most appropriate and parsimonious model, *i.e.* that possessing the fewest parameters, but which is consistent with the data. Accordingly, the selection of the most appropriate model to describe the data for each species was based on the Akaike and Bayesian information criteria, *i.e.* AIC and BIC, which were calculated as

 $AIC = -2\lambda + 2k$  $BIC = -2\lambda + k \ln n$ 

where *k* is the number of parameters, *i.e.* 2 for the linear, 3 for the quadratic, 4 for the broken stick regression and 6 for the overlapping-lines model (Akaike, 1973; Schwartz, 1978). The model with the smallest value for the information criterion was accepted as providing the best, most parsimonious fit to the data for that species and for that information criterion.

#### 7.2.4 Probability of selecting an incorrect model

The parameters that were estimated when the linear model was fitted to each set of randomly resampled data for each of *P. pelagicus*, *H. acerba* and *C. bicolor*, and the resulting standard deviation of the observation errors were then used to generate a synthetic value of the logarithm of dorsal propodus length for each recorded body size. For this, a random normal variate with a mean of zero and standard deviation of 1 was generated. This was multiplied by the standard deviation of the observation errors to produce a random "observation error", which was then added to the value predicted using the linear model and the recorded body size to produce a randomly-generated value of the logarithm of dorsal propodus length for the crab. The resulting data are thus formed from the linear model fitted to the resampled data, *i.e.* to data randomly drawn from the population from which the original data set was obtained. Linear, quadratic, broken-stick and overlapping-lines models were then fitted to the synthetic data and, for each of the AIC and BIC, the best, most parsimonious model was selected as that with the lowest value of that information criterion. This process of generating a synthetic set of data, fitting the three models and selecting the more appropriate model was repeated using each set of results obtained from refitting the linear model to each of the resampled data sets. The probability of erroneously selecting either the quadratic, broken-stick or overlapping-lines model as the best model was calculated as the proportion of these 200 runs for which each of these models was selected as the best model, *i.e.* the correct conclusion that there was no change in the level of allometry was rejected.

The same process was then repeated for each species, using the parameter estimates obtained when fitting the quadratic, then the broken-stick, and finally the overlapping lines, models to generate the random data to which the other models were fitted and to determine whether or not the model that matched that used to generate the synthetic data was correctly selected using the AIC and the BIC.

124

We have considered four hypotheses concerning the relationship between the logarithms of the length of the dorsal propodus and of body size, *i.e.*  $H_1$  - the relationship is linear,  $H_2$  - the relationship is quadratic,  $H_3$  - the relationship is a broken stick or  $H_4$  - the relationship is one that is formed by overlapping lines. The conclusion from our analysis is that most appropriate relationship for describing the data is either  $D_1$  - linear,  $D_2$  - quadratic,  $D_3$  - a broken stick, or  $D_4$  - one that is formed by overlapping lines.

The results obtained from the simulations described in the preceding paragraphs were used to calculate the probability of each conclusion, D, given that the relationship used to generate the synthetic data was known to be of each specific form. This has been denoted as  $P_{AIC}(D|H)$  or  $P_{BIC}(D|H)$ . Thus, for example,  $P_{BIC}(D = D_1 | H_3)$  is estimated as the proportion of data sets that were generated using the broken stick relationship in which, employing the BIC, it was concluded that the relationship was linear.

A Bayesian approach was employed to assign probabilities to each of the hypotheses regarding the form of the relationship between the logarithms of the length of the dorsal propodus and the body size. It was assumed that, prior to the study, the four hypotheses were equally likely, *i.e.* P(H) = 0.25 for each hypothesis, *H*. The posterior probabilities of these hypotheses were then calculated as

$$P(H_j) = \frac{P(D \mid H_j)P(H_j)}{\sum_{k} P(D \mid H_k)P(H_k)}.$$

#### 7.2.5 Physiological maturity

Collected individuals of *H. acerba* and *C. bicolor* were transported to the laboratory where they were euthanased in a bath of seawater and clove oil. The gonads of these crabs in a randomly-selected subset of the males of each species, which could be readily distinguished from females by the shape of their abdomen, were examined macroscopically and classified as immature or mature using the following criteria. They were recorded as immature if either the reproductive tract, *i.e.* the testes and/or vas deferens, was not visible or if the middle and posterior vas deferens were present but were thin and either straight or loosely convoluted, and were classified as mature when the middle and posterior regions of the vas deferens were enlarged and highly convoluted (Ryan, 1967; de Lestang *et al.*, 2003a). Note that, when examined microscopically, segments of the anterior vas deferens of large subsamples of each species always contained spermatophores in individuals designated as mature, but that this was never the case with immature crabs.

#### 7.2.6 Size at physiological maturity

The sizes by which 50% of the males of *H. acerba* and *C. bicolor* have become physiologically mature was calculated by logistic regression analysis, using the equation

$$p_{j} = \{1 + \exp[-\ln(19)(CL_{j} - CL_{50})/(CL_{95} - CL_{50})]\}^{-1},\$$

to calculate  $p_{j_i}$  the probability that a male crab of carapace width  $CL_j$  is mature and where  $CL_{50}$ and  $CL_{95}$  are the carapace widths at which 50 and 95% of the males are expected to be mature. This equation is a reparameterised version of the logistic model, in which the traditional parameters are replaced with the parameters  $CL_{50}$  and  $CL_{95}$ . The model was fitted by maximum likelihood using a nonlinear subroutine in SPSS<sup>TM</sup>.

#### 7.3 RESULTS

#### 7.3.1 Relationship between propodus length and body size.

Comparison of the values of the AIC and BIC, calculated by fitting each model to the logarithms of *DP* and *CW* for the males of *P. pelagicus*, demonstrated that the relationship

between these measures was described better by a broken stick regression than by either a

linear, quadratic or overlapping-lines model (Table 7.1, Figure 7.1).

**Table 7.1**. Estimates of the parameters (and 95% confidence limits) for the linear, quadratic, broken stick and overlapping-lines models fitted to the logarithms of the lengths of dorsal propodus versus logarithms of body sizes for *P. pelagicus*, *H. acerba* and *C. bicolor*, and values of the log-likelihood ( $\lambda$ ) and of the Akaike (AIC) and Bayesian (BIC) Information Criteria.

		P. pelagicus			H. acerba			C. bicolor		
		Estimate	Lower	Upper	Estimate	Lower	Upper	Estimate	Lower	Upper
Linear	а	-3.003	-3.103	-2.917	-3.029	-3.196	-2.884	-1.741	-1.814	-1.656
	b	1.374	1.355	1.396	1.539	1.507	1.576	1.147	1.128	1.162
	λ	2270	2228	2314	1018	988	1058	1692	1642	1750
	AIC	-4536	-4623	-4452	-2033	-2111	-1971	-3380	-3495	-3279
	BIC	-4526	-4614	-4443	-2025	-2103	-1963	-3371	-3487	-3270
Quadratic	а	0.699	-0.154	1.732	-3.619	-5.958	-1.116	0.265	-0.986	1.756
	b	-0.374	-0.858	0.018	1.807	0.693	2.854	0.285	-0.346	0.822
	С	0.205	0.161	0.259	-0.030	-0.148	0.099	0.092	0.035	0.161
	λ	2390	2340	2461	1019	989	1058	1694	1643	1755
	AIC	-4774	-4916	-4674	-2031	-2109	-1972	-3382	-3504	-3280
	BIC	-4760	-4902	-4660	-2019	-2097	-1960	-3370	-3491	-3267
Broken stick	d	-2.384	-2.559	-2.256	-3.143	-12.916	-0.060	-1.335	-1.677	5.268
	е	1.222	1.190	1.267	1.565	0.801	4.162	1.055	-0.521	1.133
	f	1.566	1.539	1.626	1.258	0.616	1.606	1.169	1.149	1.562
	X	4.382	4.344	4.479	4.705	3.879	4.804	4.511	4.205	4.991
	$\exp(X)$	80.000	77.000	88.167	110.495	48.361	122.000	91.000	67.000	147.025
	λ	2402	2356	2463	1022	992	1062	1695	1644	1758
	AIC	-4796	-4919	-4704	-2036	-2115	-1976	-3383	-3507	-3280
	BIC	-4777	-4900	-4685	-2020	-2099	-1960	-3365	-3490	-3263
Overlapping	a'	-2.403	-2.583	-2.259	-2.675	-3.136	-2.223	-1.380	-2.010	-1.110
lines	b'	1.227	1.191	1.272	1.453	1.345	1.561	1.066	1.006	1.212
	с'	-3.891	-4.156	-3.751	-2.557	-2.891	-1.427	-1.724	-2.405	-1.358
	ď	1.566	1.536	1.623	1.438	1.199	1.511	1.144	1.070	1.279
	$S_{50}$	76.535	73.221	88.610	85.891	83.743	97.876	106.093	86.467	132.911
	$S_{95}$	77.304	74.039	91.543	86.754	84.585	104.900	107.159	87.336	134.247
	λ	2402	2357	2465	1026	996	1066	1698	1647	1760
	AIC	-4792	-4918	-4701	-2041	-2121	-1979	-3384	-3508	-3282
	BIC	-4764	-4890	-4673	-2017	-2097	-1956	-3358	-3482	-3257
		BS			Overlap			Overlap		
		BS			Linear			Linear		

However, the use of the AIC suggested that, for the males of both *H. acerba* and *C. bicolor*, the overlapping-lines model provided the best description of the relationship between the



**Figure 7.1.** Alternative model forms (linear, quadratic, broken stick and overlapping lines) fitted to the relationship between the natural log of the length of the dorsal propodus of the largest cheliped and the natural log of the carapace width of *Portunus pelagicus* and the natural log of the carapace length for *Hypothalassia acerba* and *Chaceon bicolor*.

logarithms of *DP* and *CL*, whereas the use of the BIC indicated that the linear model was more appropriate (**Table 7.1**).

#### 7.3.2 The probability of selecting an incorrect model form

The percentages of synthetic data sets for which each model was correctly or incorrectly identified by the AIC or BIC as providing the best description of the relationship between the  $\ln(DP)$  and  $\ln(CW)$  or  $\ln(CL)$  are presented in **Table 7.2**. These data were then used to generate the posterior probabilities shown in **Table 7.3**, the implications of which are considered in the discussion.

#### 7.3.3 Size at physiological maturity

The estimates of  $CW_{50}$  and  $CW_{95}$  for *Portunus pelagicus*, based on the percentages of male crabs with mature gonads, were 88.4 (95% confidence interval 87.8 – 89.1 mm) and 99.9 (95% C.I. 98.2 – 101.6) mm, respectively (de Lestang *et al.*, 2003a) (**Figure 7.2**). Males of *H. acerba* attain physiological maturity at a carapace length of 68.1 (95% C.I. 67.8 - 68.3) mm, and 95% are physiologically mature when they have reached a carapace length of 72.0 (95% C.I. 71.2 – 72.7) mm. For male *Chaceon bicolor*, the estimates of  $CL_{50}$  and  $CL_{95}$  were 94.3 (95% C.I. 93.7 - 94.9) and 99.9 (95% C.I. 98.2 – 101.6) mm CL, respectively (**Figure 7.2**).

#### 7.4 DISCUSSION

#### 7.4.1 Morphometric and physiological maturity of Portunus pelagicus

Our results show that, irrespective of whether the selection of model form was based on AIC or BIC, the relationship between the logarithms of the dorsal propodus and carapace width of *P. pelagicus* was described better using the broken stick approach rather than either linear, quadratic or overlapping-lines models. Although the posterior probabilities estimated for the broken stick and overlapping-lines models for this species differed only slightly, but were both



**Figure 7.2.** Logistic regressions fitted to the proportions of males of (a) *Portunus pelagicus* (b) *Hypothalassia acerba* and (c) *Chaceon bicolor* which at each size possessed mature gonads.

**Table 2.** Percentage of data sets in which, using either the Akaike (AIC) or Bayesian (BIC) Information Criterion, each model was selected as the model of appropriate complexity that best described the 200 sets of synthetic random data generated using each of the linear, quadratic, broken stick and overlapping-lines models. The correct classification is identified by the shading.

		Model used to generate synthetic data							
	Selected	Linear		Quadratic		Broken stick		Overlap	
Species	model	AIC	BIC	AIC	BIC	AIC	BIC	AIC	BIC
P. pelagicus	Linear	64.5	98.5	0.0	0.0	0.0	0.0	0.0	0.0
	Quadratic	9.0	1.0	99.0	100.0	0.0	0.0	0.0	0.0
	Broken stick	21.0	0.5	1.0	0.0	85.5	100.0	72.0	99.5
	Overlap	5.5	0.0	0.0	0.0	14.5	0.0	28.0	0.5
H. acerba	Linear	62.5	98.5	44.0	87.0	7.0	35.5	2.0	43.0
	Quadratic	10.5	1.0	21.0	11.5	15.0	30.5	0.5	0.0
	Broken stick	19.5	0.5	25.5	1.5	69.5	33.5	11.0	10.0
	Overlap	7.5	0.0	9.5	0.0	8.5	0.5	86.5	47.0
C. bicolor	Linear	63.5	99.0	21.5	54.5	10.5	44.5	2.5	42.5
	Quadratic	7.5	0.5	39.5	43.5	22.5	38.5	8.5	29.0
	Broken stick	19.5	0.5	32.0	2.0	51.5	17.0	33.5	20.0
	Overlap	9.5	0.0	7.0	0.0	15.5	0.0	55.5	8.5

**Table 3.** Posterior probabilities of the model forms describing the allometric data for each species calculated from the fact that, using both the Akaike (AIC) and the Bayesian (BIC) Information Criterion, the broken stick model was selected as the most appropriate model to represent the allometric data for *P. pelagicus*, and that, using the AIC, the overlapping-lines model was selected as the most appropriate model to describe the data for *H. acerba* and *C. bicolor* whereas the linear model was chosen when the BIC was employed.

	Posterior probability of each model form for each species								
	P. pelagicus		H. ac	cerba	C. bicolor				
Model form	AIC	BIC	AIC	BIC	AIC	BIC			
Linear	0.117	0.003	0.067	0.373	0.109	0.412			
Quadratic	0.006	0.000	0.085	0.330	0.080	0.227			
Broken stick	0.476	0.500	0.076	0.134	0.177	0.185			
Overlapping	0.401	0.498	0.772	0.163	0.634	0.177			
lines									

higher than those of either the linear or quadratic models, the broken stick is less complex than the overlapping-lines model (four as opposed to six parameters) and is thus the better and more appropriate model.

From our use of the broken stick approach, it is concluded that the growth of the largest cheliped of male *P. pelagicus* clearly undergoes an inflection when individuals reach a carapace width of about 80.0 mm (95% confidence limits 77.0 to 88.2 mm). This value is ~ 6 mm less than that derived previously by de Lestang *et al.*, (2003a), *i.e.* 86.2 mm CW, when subjecting the same data to Somerton's (1980) overlapping-lines method. However, in this context, it is relevant that the parameter estimates of the overlapping-lines model for *P. pelagicus* have been found to be very sensitive to the values selected as the upper and lower "limits" of the body sizes associated with morphometrically immature or mature individuals. Thus, the estimate derived from the previous study is unlikely to have been precise.

The males of *P. pelagicus* become physiologically mature at 88.4 mm CW (de Lestang *et al.*, 2003a), and thus ~8 mm after they have attained morphometric maturity. Thus, the males of this species would appear, at least on the basis of cheliped measurements, to have developed the potential to mate just prior to the time when their gonads become fully mature. From a management perspective, it would be appropriate to base management strategies and assessment on the size at which *P. pelagicus* has attained both morphometric and physiological maturity, *i.e.* 88.4 mm CW.

# 7.4.2 Morphometric and physiological maturity of Hypothalassia acerba and Chaceon bicolor

The values estimated for the AIC and BIC for the males of *H. acerba* and *C. bicolor* gave no consistent indication of which of the four models provides the best description of the relationship between the logarithms of the length of the dorsal propodus and CL. For both species, the value of the AIC suggested that the overlapping-lines model was best, whereas the value of the BIC indicated that the linear model was best. Since the values of the posterior probabilities that were estimated for each model showed the same inconsistency, there is a no conspicuous change in the level of allometry of the males of either *H. acerba* or *C. bicolor*, that could be related to the attainment of morphometric maturity.

The absence of both distinct changes in the patterns of allometry of the chelae of the males of *H. acerba* and *C. bicolor* and direct information on the sizes at which these deep-sea crabs are capable of mating, *i.e.* functional maturity, leaves only the size of physiological maturity as a basis for assessing the "reproductive" status of their stocks and as a measure of reproductive potential for developing management plans for these species. Thus, since *H. acerba* and *C. bicolor* become physiologically mature at 68.1 and 94.3 mm CL, respectively, it

would be appropriate for managers to consider these lengths when reviewing management plans involving the use of a minimum legal size.

This paper has been submitted for publication as follows: Hall, N.G., Smith, K.D., de Lestang, S. and Potter, I.C. (submitted). Do the chelae of the males of three crab species undergo allometric changes that can be used to determine morphometric maturity. *ICES Journal of Marine Science*.

#### 8.0 **REFERENCES**

- Abbe, G. R. 2002. Decline in size of male blue crabs (*Callinectes sapidus*) from 1968 to 2000 near Calvert Cliffs, Maryland. Estuaries **25**: 105-114.
- Akaike, H. 1973. Information theory and an extension of the maximum likelihood principle. pp. 267-281. In B. N. Petrov and F. Csaki (eds). Second International Symposium on Information Theory. Akademiai Kiado, Budapest
- Anonymous. 2002. Western Australian Department of Fisheries, Catch and Effort Statistics.
   Fisheries Western Australia, WA Marine Research Laboratories, West Coast Drive,
   Waterman, 6020, Perth, Australia.
- Attrill, M. J., Hartnoll, R. G. and Rice, A. L. 1991. Aspects of the biology of the deep-sea crab *Geryon trispinosus* from the Porcupine Seabight. J. Mar. Biol. Ass. U.K. **71**: 311-28.
- Attrill, M. J., Rice, A. L., Thurston, M. H. and Hartnoll, R. G. 1990. A depth-related distribution of the red crab, *Geryon trispinosus* (Herbst) [=*G. tridens* Kroyer]: indications of vertical migration. Prog. Oceanogr. 24: 197-206.
- Bennett, D. B. & Brown, C. G. 1983. Crab (*Cancer pagurus*) migrations in the English Channel. J. Mar. Biol. Ass. U.K. 63: 371-398.
- Beyers, C. J. de B. and Wilkie, C. G. 1980. Quantitative stock survey and some biological and morphometric characteristics of the deep-sea red crab *Geryon quinquedens* off South West Africa. Fish. Bull. S.A. 13: 9-19.
- Bovbjerg, R. V. 1956. Some factors affecting aggressive behaviour in crayfish. Physiol. Zool.29: 127-136.
- Brown, R. B. and Powell, G. C. 1972. The size at maturity in the male Alaskan tanner crab, *Chionocetes bairdi*, as determined by chelae allometry, reproductive tract weights, and size of pre-copulatory males. J. Fish. Res. Bd Can. 29: 423–427.

- Caddy, J. F. and Mahon, R. 1995. Reference points for fisheries management. FAO Fisheries Technical paper 347. 83pp.
- Carroll, J. C. and Winn, R. N. 1989. Brown rock crab, red rock crab, and yellow crab. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southwest). Biological Report of the National Wetlands Research Centre U.S. Fish and Wildlife Service, 11: 117. 18pp.
- Cerrato, R. M. 1990. Interpretable statistical tests for growth comparisons using parameters in the von Bertalanffy equation. Can. J. Fish Aquat. Sci. **47**: 1416-1426.
- Clarke, K. R., and Gorley, R. N. (2001). PRIMER v 5: User Manual/Tutorial. Primer E: Plymouth.
- Clayton, D. A. 1990. Crustacean allometric growth: a case for caution. Crustaceana **58**: 270-290.
- Comeau, M., and Conan, G. Y. 1992. Morphometry and gonad maturity of male snow crab *Chionoecetes opilio*. Can. J. Fish. Aquat. Sci. 49: 2460-2468.
- Comeau, M., Conan, G. Y., Matnou, F., Robichaud, G., Therriault, J-C. and Starr, M. 1998.
  Growth, spatial distribution, and abundance of benthic stages of the snow crab
  (*Chionoecetes opilio*) in Bonne Bay, Newfoundland, Canada. Can. J. Fish. Aquat. Sci.
  55: 262-79.
- Comeau, M., Conan, G. Y., Maynou, F., Robichaud, G., Therriault. J-C. and Starr, M. 1998.
  Growth, spatial distribution, and abundance of benthic stages of the snow crab (*Chionoectes opilio*) in Bonne Bay, Newfoundland, Canada. Can. J. Fish. Aquat. Sci. 55: 262-279.
- Conan, G. Y. and Comeau, M. 1986. Functional maturity and terminal molt of male snow crab, *Chionoecetes opilio*. Can. J. Fish. Aquat. Sci. **43**: 1710-1719.

- Conan, G. Y., Comeau, M. and Moriyasu, M. 2001. Are morphometrical approaches appropriate to establish size at maturity for male American lobster, *Homarus americanus*? J. Crust. Biol. 21: 937-947.
- Defeo, O., Little, V. and Barea, L. 1991. Stock assessment of the deep-sea red crab *Chaceon notialis* in the Argentinian-Uruguayan common fishing zone. Fish Res. **11**: 25-39.
- de Lestang, S., Hall, N. G. and Potter, I. C. 2003a. Reproductive biology of the blue swimmer crab (*Portunus pelagicus*, Decapoda: Portunidae) in five bodies of water on the west coast of Australia. Fish. Bull US **101**: 745-757.
- de Lestang, S., Hall, N., and Potter, I. C. 2003b. Influence of a deep artificial entrance channel on the biological characteristics of the blue swimmer crab *Portunus pelagicus* in a large microtidal estuary. J. Exp. Mar. Biol. Ecol. **295**: 41-61.
- Ennis, G. P., Hooper, R. G. and Taylor, D. M. 1988. Functional maturity in small male snow crabs (*Chionoecetes opilio*). Can. J. Fish. Aquat. Sci. **45**: 2106-2109.
- Fernández-Vergaz, V., López Abellán, L. J. and Balguerías, E. 2000. Morphometric, functional and sexual maturity of the deep-sea red crab *Chaceon affinis* inhabiting Canary Island waters: chronology of maturation. Mar. Ecol. Prog. Ser. **204**: 169-178.
- Fisher, M. R. 1999. Effect of temperature and salinity on size at maturity of female blue crabs. Tafs Trans. Am. Fish. Soc. 128: 499-506.
- Fournier, D. A. 2000. An Introduction to AD Model Builder Version 4: For Use in Nonlinear Modelling and Statistics. Otter Research, Sidney, Canada. 127pp.

Gardner, C. 1996. Behavioral basis of depth regulation in the first zoeal stage of the giant crab (*Pseudocarcinus gigas*, Brachyura, Xanthoidea, Oziidae). In High latitude crabs: biology, management, and economics. pp. 229-54. University of Alaska Sea Grant College Program Report No. 96-02.

- Gardner, C. 1997. Effect of size on reproductive output of giant crabs *Pseudocarcinus gigas* (Lamarck): Oziidae. Mar. Freshwat. Res. **48**: 581-587.
- Gardner, C. 1998a. The first record of larvae of the giant crab *Pseudocarcinus gigas* in the plankton. Pap. Proc. R. Soc. Tas. **132**: 47-48.
- Gardner, C. 1998b. *The Tasmanian giant crab fishery: A synopsis of biological and fisheries information*. Department of Primary Industry and Fisheries, Marine Division: Tasmania.
- Gardner, C. and H. Williams. 2002. Maturation in male giant crab, *Pseudocarcinus gigas*, and the potential for sperm limitation in the Tasmanian fishery. Mar. Freshwat. Res. **53**: 661-667.
- George, R. W. 1962. Description of *Panulirus cygnus* sp. nov., the commercial crayfish (or spiny lobster) of Western Australia. J. Roy. Soc., WA 45, 100-110.
- George, R. W. 1966. *Hypothalassia armata* (De Haan) in Western Australia. Crustaceana **10**, 223-224.
- Goshima, S. and Kanazawa, M. 2000. Maturity in male snow crab *Haplogaster dentata* (Anomura: Lithodidae) and its application for fishery management. J. Crust. Biol. 20: 641-646.
- Goshima, S., Kanazawa, M., Yoshino, K. and Wada, S. 2000. Maturity in male stone crab *Haplogaster dentata* (Anomura: Lithodidae) and its application for fishery management.
  J. Crust. Biol. 20: 641-646.
- Groeneveld, J. C. and Branch, G. M. 2002. Long-distance migration of South African deepwater rock lobster *Palinurus gilchristi*. Mar. Ecol. Prog. Ser. **232**: 225-238.
- Haefner, P. 1977. Reproductive biology of the female deep sea red crab, *Geryon quinquedens*, from the Chesapeake Bight. Fish. Bull. **75**: 91-100.

- Haefner, P. A. 1978. Seasonal aspects of the biology, distribution and relative abundance of the deep-sea red crab *Geryon quinquedens* Smith, in the vicinity of the Norfolk Canyon, western North Atlantic. Proc. Nat. Shellfish Ass. 68: 49-62.
- Hartnoll, R. G. 1978. The determination of relative growth in Crustacea. Crustaceana **34**: 281-293.
- Hastie, L. C. 1995. Deep-water geryonid crabs: a continental slope resource. Oceanogr. Mar. Biol. Ann. Rev. 33: 561-584.
- Hooper, R. G. 1986. A spring breeding migration of the snow crab, *Chionoecetes opilio* (O. Fabr.), into shallow water in Newfoundland. Crustaceana 50, 257-64.
- Ingles, J. and Braum, E. 1989. Reproduction and larval ecology of the blue swimming crab *Portunus pelagicus* in Rabay Gulf, Phillipines. International Revue der gesamten Hydrobiologie **74**: 471-490.
- Jivoff, P. and Hines, A. H. 1998. Female behaviour, sexual competition and mate guarding in the blue crab, *Callinectes sapidus*. Anim. Behav. 55: 589-603.
- Jones, D. S. and Morgan, G J. 2002. *A field guide to crustaceans of Australian waters*. New Holland Publishers, Sydney.
- Jury, S. H. and Howell, H. 2001. Lobster trap video: in situ video surveillance of the behaviour of *Homarus americanus* in and around traps. Mar. Freshwat. Res. **52**: 1125-32.
- Kailola, P. J., Williams, M. J., Stewart, P. C., Reichelt, R. E., McNee, A. and Grieve, C. 1993.
  Australian Fisheries Resources. Canberra: Department of Primary Industry and Energy.
  422 pp.
- Koh, S. K. and Ng P. K. L. 2000. A revision of the spiny crabs of the genus *Hypothalassia* Gistel, 1848 (Crustacea: Decapoda: Brachyura: Eriphidae). Raffles Bull. Zool. 40: 123-41.

- Levings, A., Mitchell, B. D., Heeren, T. and Austin, C. 1996. Fisheries biology of the giant crab (*Pseudocarcinus gigas*, Brachyura, Oziidae) in Southern Australia. pp. 125-152. In High Latitude Crabs: Biology, Management, and Economics. Proceedings of the International Symposium on Biology, Management, and Economics of Crabs from High Latitude Habitats. University of Alaska Sea Grant College Program Report No. 96-02.
- Levings, A., B., Mitchell, D., McGarvey, R., Mathews, J., Laurenson, L., Austin, C., Heeren, T., Murphy, N., Miller, A., Rowsell, M. and Jones, P. 2001. Fisheries biology of the giant crab *Pseudocarcinus gigas*. 388 pp. Final Report to the Fisheries Research Development Corporation for Projects 93/220 and 97/132. Fisheries Research and Development Corporation, Canberra.
- Lockhart, F. D., Lindberg, W. J., Blake, N. J., Erdman, R. B., Perry, H. M. and Waller, R. S. 1990. Distributional differences and population similarities for two deep sea crabs (Family Geryonidae) in the northeastern Gulf of Mexico. Can. J. Fish. Aquat. Sci. 47: 2112-2122.
- López Abellán, L. J., Balguerias, E. and Fernandez-Vergaz, V. 2002. Life history characteristics of the deep-sea crab *Chaceon affins* population off Tenerife (Canary Islands). Fish. Res. **58**: 231-239.
- Lux, F. E., Ganz, A. R. and Rathjen, W. F. 1982. Marking studies on the red crab *Geryon quinquedens* Smith off southern New England. J. Shellfish Res. **2**: 71-80.
- McGarvey, R., Levings, A. H. and Mathews, J. M. 2002. Moulting growth of the Australian giant crab, *Pseudocarcinus gigas*. Mar Freshwat Res **53**: 869-881.
- Melville Smith, R. 1986. Red crab *Geryon maritae* density in 1985 by technique of effective area fished per trap on the northern fishing grounds off South West Africa. S. Afr. J. Mar. Sci. 4: 257-63.

- Melville Smith, R. 1987a. The reproductive biology of *Geryon maritae* (Decapoda, Brachyura) off South West Africa/Namibia. Crustaceana **53**: 11-27.
- Melville Smith, R. 1987b. Movements of deep-sea red crab (*Geryon maritae*) off South West Africa/Namibia. S. Afr. J. Zool. **22**: 143-152.
- Melville Smith, R. 1988. The commercial fishery for and population dynamics of red crab *Geryon maritae* off South West Africa. S. Afr. J. Mar. Sci. **6**: 79-95.
- Melville Smith, R. and Bailey, G. W. 1989. A preliminary investigation into the possible causes of depth zonation by red crab (*Geryon maritae*), off Namibia. Selected Papers of the International Commission for the southeast Atlantic Fisheries 1: 23-34.
- Melville Smith, R. and Cheng, Y. W. 2002. Responses of migrating western rock lobsters *Panulirus cygnus* (George, 1962) to two different tagging release procedures. J. Shellfish Res. 21: 289-294.
- Melville Smith, R., Gould, R. and Bellchambers, L. in press. The crystal crab fishery inWestern Australia: First steps in the development of a sustainable deep-water crabfishery. In Proceedings of the Deep-sea 2003 Conference. Wellington, New Zealand.
- Miller, R. J. 1990. Effectiveness of crab and lobster traps. Can. J. Fish. Aquat. Sci. **47:** 1228-1251.
- Moore, R. and MacFarlane, J. W. 1984. Migration of ornate rock lobster, *Panulirus ornatus* (Fabricus), in Papua New Guinea. Aust. J. Mar. Freshwat. Res. **35**: 197-212.
- Muino, R., Fernandez, L., Gonzalez-Gurriaran, E., Freire, J. and Vilar, J. A. 1999. Size at maturity of *Liocarcinus depurator* (Brachyura: Portunidae) a reproductive and morphometric study. Mar. Biol. **79**: 295-303.
- Paul, A. J. 1992. A review of size at maturity in male tanner (*Chionoecetes bairdi*) and king (*Paralithodes camtschaticus*) crabs and the methods used to determine maturity. Am. Zool. **32**: 534-540.

- Pearce, A. E. and Griffiths, R. W. 1991. The mesoscale structure of the Leeuwin Current: a comparison of laboratory models and satellite imagery. J. Geophys. Res. 96: 16739-16757.
- Phillips, B. F. 1983. Migrations of pre-adult western rock lobsters, *Panulirus cygnus*, in Western Australia. Mar. Biol. **76**: 311-318.
- Phillips, B. F., Brown, P. A., Rimmer, D. W. and Reid, D. D. 1979. Distribution and dispersal of the phyllosoma larvae of the western rock lobster, *Panulirus cygnus*, in the southeastern Indian Ocean. Aust. J. Mar. Freshwat. Res. **30**: 773-783.
- Phillips, B. F., Pearce, A. F., Litchfield, R. T., Pearce, A. F. and Walker, D. I. 1991. The Leeuwin Current and larval recruitment to the rock (spiny) lobster fishery off Western Australia. J. Roy. Soc. West. Aust. 74: 93-100.
- Potter, I. C., and de Lestang, S. 2000. Blue swimmer crab *Portunus pelagicus* in Leschenault Estuary and Koombana Bay, south-western Australia. J. Roy. Soc. West. Aust. **83**: 221-236.
- Rodhouse, D. M. 1984. Experimental fishing for the spider crab, *Maia squinado*: Sea and laboratory trials. J. Mar. Biol. Ass. U.K. 64: 251-259.
- Ryan, E. P. 1967. Structure and function of the reproductive system of the crab *Portunus* sauguinolentus (Herbst) (Brachyura: Portunidae). I The male system. Mar. Biol. Ass. India, Symp. Ser. 2: 506-521.
- Ryan, E. P. 1967. Structure and function of the reproductive system of the crab *Portunus* sauguinolentus (Herbst) (Brachyura: Portunidae). II The female system. Mar. Biol. Ass. India, Symp. Ser. 2: pp. 522-544.
- Sainte-Marie, B., Raymond, S. and Brêthes, J.-C. 1995. Growth and maturation of the benthic stages of male snow crab, *Chionoecetes opilio* (Brachyura: Majidae). Can. J. Fish. Aquat. Sci. 52: 903-924.

- Salmon, M. 1983. Courtship, mating systems, and sexual selection in decapods. pp. 143-169.In D. W. Dunham (ed.) Studies in Adaptation: The Behaviour of higher Crustacea. John Wiley and Sons, New York.
- Schwartz, G. 1978. Estimating the dimensions of a model. Ann. Stat. 6: 461–464.
- Somerton, D. A. 1980. A computer technique for estimating the size of sexual maturity in crabs. Can. J. Fish. Aquat. Sci. 37: 1488-1494.
- Somerton, D. A. 1981. Regional variation in the size of maturity of two species of tanner crab (*Chionoecetes bairdi* and *C. opilio*) in the eastern Bering Sea, and its use in defining management subareas. Can. J. Fish. Aquat. Sci. 38: 163-174.
- Stewart, J. and Kennelly, S. J. 1998. Contrasting movements of two exploited Scyllarid lobsters of the genus *Ibacus* off the east coast of Australia. Fish. Res. 36: 127-132.
- Tyler, P. A. 1988. Seasonalilty in the deep sea. Oceanog. Mar. Biol. Ann. Rev. 26: 227-258.
- Van Engel, W. A. 1958. The blue crab and its fishery in Chesapeake Bay. Part 1 Reproduction, early development, growth and migration. Comm. Fish. Rev. **20**: 6-17.

Warner, G. F. (ed.) 1977. The Biology of Crabs. Elek Science, London. 202 pp.

- Watson, J. 1970. Maturity, mating and egg laying in the spider crab, *Chionocetes opilio*. J. Fish. Res. Bd Can. 27: 1607-1616.
- Wigley, R. L., Theroux, R. B. and Murray, H. E. 1975. Deep-sea red crab, *Geryon quinquedens*, survey off northeastern United States. Mar. Fish. Rev. 37: 1-21.
- Williams, M. J. and Hill, B. J. 1982. Factors influencing pot catches and population estimates of the Portunid crab *Scylla serrata*. Mar. Biol. **71**: 187-192.
- Zhou, S. and Shirley, T. C. 1997. Performance of two red king crab pot designs. Can. J. Fish. Aquat. Sci. **54**: 1858-1864.

#### 9.0 **BENEFITS**

Managers now have the reliable data on the following which will enable them to produce much more effective and appropriate methods for managing the resources for the deep sea crabs *Hypothalassia acerba* and *Chaceon bicolor* (see 11.0 Planned outcomes).

- 1. Trends exhibited by annual and monthly catches, effort and catch per unit effort for *H*. *acerba* and *C. bicolor* on both the lower west and south coasts of Western Australia.
- 2. The distribution of *H. acerba* and *C. bicolor* by water depth and temperature.
- 3. The reproductive biology of *H. acerba* and *C. bicolor*.
- 4. Demonstration that some traditional approaches for estimating the size at maturity of *H*. *acerba* and *C. bicolor* are inappropriate and this may be the case for some other crustacean species.

The data will also enable fishers to make more informed decisions regarding investment in fishing for *H. acerba* and *C. bicolor*.

#### **10.0 FURTHER DEVELOPMENT**

Once the final report has been accepted by the FRDC, it will be provided to the Western Australian Department of Fisheries. Thus, when developing management plans for *H. acerba* and *C. bicolor*, fisheries managers will be able to take into account the annual and monthly trends exhibited by the fisheries catch and effort statistics and the quantitative information on key aspects of the biology of these two deep sea crab species. The final report will also be provided to WAFIC and professional fishers for deep sea crabs.

A poster will be produced describing the main findings of this study. An account of these findings will also be supplied to PROWEST so that professional fishers are made aware of the results and their implications.
## **11.0 PLANNED OUTCOMES**

The implications of our results will require managers to consider the following when they are reviewing plans for the management of *H. acerba* and *C. bicolor*.

- Estimates of the size at maturity of males should be based on physiological (gonadal) maturity.
- 2. Estimates of the size at physiological (gonadal) maturity of females are likely to be artificially low.
- 3. Values selected as the minimum legal sizes for retention of *H. acerba* and *C. bicolor* are likely to be based on the sizes at physiological maturity of males and females of these species. However, since estimates of the sizes at physiological maturity of males and females are likely to be low, an adaptive approach will be required to ensure that the numbers of mature crabs are maintained at a level sufficient to sustain the stocks.
- 4. Protection should be considered for the assemblage of *H. acerba* on the south coast as there is strong evidence that females in these waters do not typically reach maturity and apparently move to the west coast for breeding.
- 5. Restriction of both *H. acerba* and *C. bicolor* to a narrow depth range makes these species particularly subject to depletion through focussed fishing.
- 6. As the *H. acerba* and *C. bicolor* that are caught in traps appear to have approximately attained their asymptotic body size, yield per recruit will be an asymptotic function of fishing effort (of a form determined by the Baranov catch equation). Similarly, spawning biomass per recruit will decline exponentially with fishing effort. Since natural mortality cannot be estimated, it is impossible to estimate the value of fishing mortality from the available data. However, the implications of declining spawning biomass per recruit with fishing effort is that catch per unit of effort (CPUE) will need to be sustained above some reference value determined from either the virgin CPUE or

from historical levels of CPUE. Thus, decision rules for the fishery will need to be based on CPUE rather than being model-derived estimates of fishing mortality or biomass.

- If CPUE is to be used as the primary indicator for fisheries management decisions, factors that increase the efficiency of fishing effort will need to be carefully monitored. The potential for spatial depletion also exists and CPUE will need to be estimated appropriately.
- 8. Although there are no data on growth or mortality, evidence from other deep sea crab species suggest that *H. acerba* and *C. bicolor* are likely to be slow growing and long-lived. Thus, minimum legal size regulations are very likely to be inadequate unless they are accompanied by management strategies that ensure that the stocks of mature crabs are sustained at adequate levels. Since such slow growing and long-lived species are likely to be prone to becoming part of boom and bust fisheries, there is a need for constant monitoring to ascertain whether, at any stage, their catch statistics are deviating from those recorded during more recent years and thus provide evidence that the abundance of the stocks are declining. There is a strong need to consider the specification of appropriate limit and target reference points for the estimates of catch per unit of effort and for the implementation of appropriate decision rules, based on current estimates of catch per unit of effort, that will ensure that the stocks of *H. acerba* and *C. bicolor* are conserved. The decision rules should ensure that, as catch per unit of effort approaches the limit reference point for this variable, exploitation is reduced appropriately.

## **12.0 GENERAL CONCLUSIONS**

We have met the vast majority of the objectives listed in the original applications for FRDC 2001/055 and those that were our obligations for FRDC 2001/055. Please note that work associated with objectives 1, 4, 7, 8, 9 and 10 of the second proposal (FRDC 2001/055) is being undertaken by the Western Australian Department of Fisheries. The results of that work will be produced by that department in a separate final report after that project has been completed on 30 June 2005.

The only objectives we were unable to meet in FRDC 2001/055, which focused on the champagne crab *Hypothalassia acerba*, were Objectives 2 on growth and 7 on diets. Because small crabs were unable to be caught using traps, there was insufficient information in the length composition data to allow determination of the growth of individuals of either species. Without such life history data, no estimates are possible of the longevity of the individuals of either species and hence of their natural mortality. As the length composition data reveal no change that can be related to growth, yield per recruit of each crab species will increase asymptotically towards the average body weight of the individuals caught in the traps as fishing effort increases. Since the females that are caught in the traps are likely to represent the mature individuals of the stocks, the spawning female biomass per recruit of each species of crab will decrease exponentially with increasing fishing effort. The rate of increase of yield per recruit or decrease of biomass per recruit will be determined by the level of fishing mortality relative to the level of natural mortality. However, neither of these mortalities can be determined from data that can currently be collected for these species.

The diets could not be elucidated because the dietary items in the stomachs had become too decomposed for identification by the time they were examined in the laboratory. All of the objectives of FRDC 2001/055, which focused on the champagne crab *Chaceon bicolor*, were achieved. Attention is drawn to the fact that we have demonstrated that, for both *H. acerba* 

and *C. bicolor*, the traditional method for determining the size of males at morphometric maturity is not valid. This raises the strong possibility that, in the case of some other species, inappropriate methods may have been used to determine this important variable, which is widely used by fisheries managers when developing management plans. Likewise, we have demonstrated that the use of trap samples leads to underestimates of the size at maturity of *H. acerba* and *C. bicolor*. Although far better estimates can be obtained for the size of female portunids at maturity by using data from active methods of fishing, *e.g.* seine netting and trawling, these methods were not available for collecting *H. acerba* and *C. bicolor* from the deep waters in which these crabs live.

During this study, we have described the catch, effort and catch per unit for *H. acerba* and *C. bicolor* since the inception of the fisheries for those species on the lower west and south coasts of Western Australia. We have also studied the biology of those species, placing particular emphasis on determining their distribution and the factors influencing their distribution, and (2) the key aspects of the reproductive biology as they relate to management. The major findings are as follows.

- The fishery for *Hypothalassia acerba* in south-western Australia commenced in the early 1990s and peaked in 1998, with catches of ~73 000 kg. The catches subsequently declined due to fishers on the lower west coast switching their effort to catching *Chaceon acerba*. A decline in effort for *H. acerba* on the lower west coast was partially compensated for by an increase in effort on the south coast.
- The catch per unit effort for *H. acerba* has declined in recent years, suggesting that fishing might have had a detrimental influence on the stocks of this species.
- The catches of *H. acerba* on the south coast underwent a cyclical change during the year, declining to minimal levels between late autumn and mid-spring, largely

reflecting a decline in effort due to the harsh weather conditions that prevail in that period.

- The catch per unit effort of *H. acerba* on the lower west coast rose progressively from April to November, suggesting that this species gradually became more abundant and/or accessible to fishing in that period.
- The fishery for *C. bicolor*, which has been almost entirely based on the lower west coast, essentially commenced in 1997. Catches subsequently rose to reach a peack of 222 500 kg in 2000 and remained close to this level in the ensuing years.
- The sharp rise in the annual CPUE during the early years of the fishery probably reflects an increasing knowledge by fishers of the best areas to fish.
- The commercial catches of the males of both *H. acerba* and *C. bicolor* outnumbered those of their females.
- The size of the males caught by traps was greater than that of females in the case of both *H. acerba* and even more particularly *C. bicolor*.
- Traps did not catch many individuals of either *H. acerba* or *C. bicolor* which had a carapace length <80 mm, even when the traps were covered in a fine mesh.
- The catch rates of *Hypothalassia acerba* peaked in water depths of 200 m on the lower west coast and at 145 m on the south coast and in temperatures of ~16-17°C on both coasts.
- The size (carapace length) of *H. acerba* decreased slightly with increasing water depth.
- *Hypothalassia acerba* reproduces seasonally on the lower west coast, with egg extrusion occurring mainly in summer and early autumn. Since *H. acerba* does not typically appear to reproduce on the south coast, it is proposed that, maturing females migrate to the lower west coast.

- In contrast to *H. acerba*, *C. bicolor* reproduces throughout at least most of the year. This continuity of reproduction probably reflects the occupancy by this species of deeper waters than *H. acerba* and thus an environment which undergoes less conspicuous seasonal changes.
- Although the mean body weights of ovigerous females of *H. acerba* (587 g) and *C. bicolor* (652 g) were not significantly different (P > 0.05), the fecundities of these two species, *i.e.* 356, 210 and 192, 979, respectively, were significantly different. The relatively high fecundity of *H. acerba* may reflect an adaptation of this seasonal spawning species to optimise egg production.
- The passive sampling method of trapping was the only method available for catching *H. acerba* and *C. bicolor*. We thus used data collected for *Portunus pelagicus* using both the above passive fishing method and the active fishing methods of seine netting and trawling to determine whether they yielded comparable data for estimating the size at maturity of female crabs. The trap caught samples contained predominantly mature crabs and were thus biased, leading to an underestimate of the size at maturity. It is reasonable to assume that the estimates of maturity of *H. acerba* and *C. bicolor* using data derived from trap-caught samples are likewise underestimates. This point will need to be taken into account by managers when they are developing management plans for *H. acerba* and *C. bicolor*.
- We have demonstrated that the chelae of the males of *H. acerba* and *C. bicolor* do not undergo a detectable change in allometric growth at the pubertal moult. Thus, it is inappropriate to use the traditional method, which is based on this assumption, to determine the size at morphological maturity. Managers will thus have to use the size at 50% of females when they attain physiological maturity, *i.e.* 68.1 and 94.3 mm carapace lengths, respectively.

## **13.0 APPENDICES**

Appendix I

Intellectual Property

The value of the intellectual property will be 54.38% based on PART C of the FRDC project proposals.

STAFF

Professor Ian Potter Associate Professor Norm Hall Dr Alex Hesp Dr Simon de Lestang